



Review Article

ISSN : 2277-3657  
CODEN(USA) : IJPRPM

## *Urinary Complications among Women with Cystoceles: A Systematic Review*

Hatim T. O. Ali<sup>\*1</sup>, Saeed Abdullah Saeed Alqahtani<sup>2</sup>, Mohammed Saad Salem Alqahtani<sup>2</sup>, Safiyh Ahmed Alasiri<sup>2</sup>, Afnan Saeed Mgbel<sup>2</sup>, Faris Ahmed A Alasmre<sup>3</sup>, Hossien Saad Salem Alqahtani<sup>3</sup>, Ahmed Saeed S Alqahtani<sup>3</sup>, Saeed Ali S Algarni<sup>3</sup>, Abdulmajeed Samry Hassan Alanazi<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, College of Medicine, King Khalid University, Abha, Saudi Arabia.

<sup>2</sup>Department of Obstetrics and Gynecology, Maternity and Children's Hospital, Abha, Saudi Arabia.

<sup>3</sup>College of Medicine, King Khalid University, Abha, Saudi Arabia.

\*Email: [hatimtag@gmail.com](mailto:hatimtag@gmail.com)

---

### ABSTRACT

*Cystoceles result from a pelvic-floor support system that is not strong enough. This systematic review aims to study the urinary complications in women with cystoceles, especially SUI, and the complication following surgical repair. PubMed, Web of Science, Science Direct, EBSCO, and Scopus were searched. Study articles were screened by title and abstract using Rayyan QCRI then a full-text assessment was implemented. Thirteen studies were included, with a total of 1043 women with cystoceles and urinary complications. The follow-up duration ranged from 1.4 months to 53 months. Most of the included studies had SUI as a complication of cystoceles. The other complication included urge incontinence, mixed incontinence, frequency, incomplete voiding, and voiding difficulties. Incontinence recurrence and voiding dysfunction were reported only in one investigation. The majority of cystoceles cases had SUI. Frequency, incomplete voiding, urge incontinence, mixed incontinence, and voiding issues were among the other complications. The most frequent surgical complications were pyelonephritis, cystitis, and UTIs.*

**Key words:** *Cystocele, Pelvic prolapse, Urinary incontinence, Systematic review*

---

### INTRODUCTION

A cystocele often referred to as a protrusion of the bladder develops when the bladder descends into the vagina. The bladder protrudes through the anterior wall of the vagina, with which it is anatomically connected. Several underlying factors might lead to cystocele development, which causes the muscles and connective tissue surrounding the bladder and vagina to weaken [1].

In women, the endopelvic fascia and pelvic floor muscles support the pelvic viscera anatomically. Pubococcygeus, puborectalis, and iliococcygeus, three muscles that together make up the levator ani muscle group, provide this pelvic support. Additionally, the cardinal and uterosacral ligaments offer extra stability and support. The functionality of daily life and sexual activities might be negatively impacted by a prolapsed or herniated bladder. Some women may experience modest symptoms that do not need treatment, and in some circumstances, prolapses may regress after menopause [2].

Cystoceles are the result of a pelvic-floor support system that is not strong enough. Obesity, advancing age, and parity are the primary risk factors. They may also develop as a result of pelvic surgery, collagen abnormalities, a familial history of cystocele, and chronically elevated intra-abdominal pressure [2].

The exact number of women who suffer from anterior vaginal wall prolapse is difficult to determine. According to the Women's Health Initiative study, 32.9% of women who have had a hysterectomy and 34.3% of women with uteruses experience bladder prolapse [3]. 2.9% of non-pregnant women over 20 who were not pregnant were found to have some degree of pelvic organ prolapse in a cross-sectional survey of the participants [4]. The severity of the symptoms was shown to be higher in groups of women from lower socioeconomic backgrounds. Additionally, it has been determined that by age 80, 11.1% of women will need an operation to control a prolapse, with around one-third of them requiring a second surgery [5].

Only women who report symptomatic anterior wall prolapse are advised to have treatment for cystocele [6]. Generally speaking, there are three treatment options for the condition: expectant, conservative, and surgical. As part of a therapy strategy, the patient should be presented with all available options and adequately briefed on them. Better results are observed if surgical goals and the patient's expectations are discussed before treatment [7].

The next option for symptomatic women, those who need quick relief, those who have rejected conservative treatment, or for whom the conservative strategy has not been successful is surgical management. It should be highlighted that surgery should be performed after the patient's family is complete because childbirth can further harm the pelvic support system [8]. This systematic review investigates the urinary complications associated with cystoceles, especially stress urinary incontinence (SUI), and following surgical repair among women [9-11].

## MATERIALS AND METHODS

This systematic review was conducted following the accepted standards (Preferred Reporting Items for Systematic Reviews and Meta-Analyses, PRISMA).

### *Study design*

This was a systematic Review.

### *Study condition*

This study reviewed the urinary complications in women with cystoceles, especially SUI, and the complication following surgical repair.

### *Search strategy*

To find the appropriate literature, a thorough literature search was carried out in five major databases: PubMed, Web of Science, Science Direct, EBSCO, and Scopus. The English language was our only option, and the requirements of each database were taken into consideration. The following keywords, which were converted into Mesh terms in PubMed, were used to find the right studies; "Cystocele," "Pelvic prolapse," "Vaginal prolapse," "Complications," "Sequelae," "Urinary incontinence," "Urinary stress incontinence," "Urinary urge incontinence," and "Urinary tract disorders." The appropriate keywords were paired with the Boolean operators "OR" and "AND." The search yielded English-language publications with full text, freely accessible articles, and human trials.

### *Selection criteria*

#### *Inclusion criteria*

The subjects were chosen for addition founded on their applicability to the research, which has the following criteria; only female patients with cystoceles who had urinary complications.

#### *Exclusion criteria*

We dismissed all subsequent publications, ongoing research, and analyses of completed studies that did not focus on our main object.

#### *Data extraction*

To check for duplicate search strategy results, we used Rayyan (QCRI) [12]. By imposing a set of inclusion/exclusion criteria on the combined search results, the researchers were able to determine the relevance of the titles and abstracts. The reviewers read the entirety of the papers' texts that met the criteria for inclusion. The authors discussed dispute resolution. The qualified study was incorporated using a created data extraction

form. The authors extracted data about the study titles, authors, study year, study designs, participant number, mean age, follow-up duration, preoperative complications of cystoceles, type of procedure, and postoperative complications.


#### Strategy for data synthesis

To provide a qualitative summary of the included study components and outcomes, summary tables were created using the data collected from the eligible studies. After data extraction for the systematic review, the best use of the data from the included study articles was decided. Studies that met the full-text inclusion criteria but did not provide any data on urinary complications among women with cystoceles were excluded.

## RESULTS AND DISCUSSION

#### Search results

A total of 420 study articles resulted from the systematic search, and then 40 duplicates were removed. Title and abstract screening were conducted on 380 studies, and 309 studies were excluded. 71 reports were sought for retrieval, and only 3 articles were not retrieved. Finally, 68 studies were screened for full-text assessment; 29 were excluded for wrong study outcomes, 17 for unavailable data on the urinary complications of cystoceles, and 9 for the wrong population type. Thirteen eligible study articles were included in this systematic review. **Figure 1** displays a summary of the study selection method.



**Figure 1.** Summarizes the study selection procedure using a PRISMA flowchart.

#### Characteristics of the included studies

A total of 13 studies were included in this review, with 1043 patients with cystoceles and urinary complications. Five studies were retrospective [13-17], three were randomized control trials (RCTs) studies [18-20], one was a case-control study [21], one was a descriptive study [22], and one was a prospective study [23]. Three studies

were conducted in Italy [15, 20, 23], One in India [22], one in Australia [13], one in Japan [18], one in Egypt [19], one in South Korea [14], one in France [20], one in Taiwan [16], one in Georgia [24], one in Turkey [17], and one in the USA [23]. Most of the included populations were elderly. The follow-up duration ranged from 1.4 months [13] to 53 months [15]. Most of the included studies had SUI as a complication of cystoceles [13-17, 19, 20, 22-25]. The other complication included urge incontinence, mixed incontinence, frequency, incomplete voiding, and voiding difficulties. Cystocele repair was done through various procedure types. Most cases with SUI were cured following the repair. Urinary tract infection (UTI), cystitis, and pyelonephritis were reported in four studies as postoperative complications [14, 15, 20, 22]. Incontinence recurrence and voiding dysfunction were reported in one investigation [24].

**Table 1.** A summary of characteristics of the included study articles.

Study	Country	Study designs	Participants (n)	Mean age	Follow-up duration (months)	Preoperative complications	Procedure type	Postoperative complications
Shrestha, 2016 [22]	India	Descriptive study	80	15-45	NA	23.8% have SUI, and 15% have urge incontinence	Various procedures for cystocele repair	UTI followed by retention of urine
Shek et al., 2009 [13]	Australia	Retrospective study	93	61 ± 11.5	1.4-24	25.8% have SUI	Transobturator mesh	No postoperative complications. Net curative effect.
Okui et al., 2009 [21]	Japan	Case-control	34	42-89	12-37	Urgency, daytime frequency, and incontinence	Polypropylene mesh	Improvement in the preoperative symptoms; however, no significant improvement in nocturia was found.
Abdel Aziz et al., 2020 [18]	Egypt	Prospective randomized trial	72	52 ± 9.4	12	Frequency followed by urgency, urgency incontinence, weak flow, and interrupted flow.	Trans-obturator tape (TOT) and TOT with concomitant repair of cystocele by anterior colporrhaphy	The cure rate of SUI increases and the frequency of irritative urinary symptoms decrease when stage II asymptomatic cystocele and a mid-urethral sling are simultaneously repaired.
Park et al., 2013 [19]	South Korea	RCT	91	58 ± 9.5	12	43.3% have SUI, and 48.4 have mixed incontinence	Tension-free vaginal tape (TVT) and concomitant cystocele repair	Mixed incontinence was the most prevalent
Eboue et al., 2010 [14]	France	Retrospective study	123	62	15	57% have SUI, 40% with urgency, and 39% with incomplete voiding	Transobturator four arms mesh	Incomplete voiding, UTI, pyelonephritis, bladder obstruction, and de novo stress incontinence.
Zullo et al., 2008 [20]	Italy	RCT	69	49-75	12	All patients are with SUI	Anterior colporrhaphy plus inside-out TVT	Fever, UTI, severe pain, and transitory urinary retention

Cugudda et al., 2002 [15]	Italy	Retrospective study	36	57.3 ± 9.6	53	All patients are with SUI	Anterior colpoperineorrhaphy and burch colposuspension	Vaginal candidiasis and cystitis
Chen et al., 2011 [16]	Taiwan	Retrospective study	50	65.7 ± 8.9	24	32% with frequency, 28% with urinary incontinence, 6% with urgency, 10% with dysuria, 20% with voiding difficulty, and 14% with a feeling of incomplete voiding	Isolated cystocele repair	Only one patient continued to have incontinence
Bruce et al., 1999 [24]	Georgia	NA	52	61	17	All patients are with SUI	Paravaginal defect repair (PVdR)	Incontinence recurrence, voiding dysfunction, or the appearance of recurrent vaginal prolapse were late postoperative problems.
Onarimmun, 2014 [17]	Turkey	Retrospective study	165	56.3 ± 11.2	45.1±19.9	All patients are with SUI	TOT	There is a significant improvement in the SUI after cystocele repair
Gillera et al., 2006 [25]	USA	NA	121	67	NA	6% with SUI	Vaginal gauze pack	The Valsalva leak point pressures in those with SUI were indicative of intrinsic sphincter insufficiency
Adile et al., 2012 [23]	Italy	prospective study	57	57	36	20% with SUI	Monoprosthesis	80% of patients were cured of SUI

We believe that this is the first systematic review to investigate the urinary complications associated with cystoceles among women in addition to the postoperative complications. This review is mainly limited by the lack of quantitative assessment to evaluate our objective. We have fully outlined the studies we have included and have obtained data on several complications.

Another important drawback is the inclusion of low-quality evidence: most studies lacked a comparative design, did not look at treatments based on complication type, and did not compare results based on various management approaches. Therefore, it is challenging to interpret the evidence. Most studies failed to report outcomes per the reason for referral, which is crucial information for figuring out how to manage complications effectively.

The present study found that most cystoceles cases suffered from USI. Urge incontinence, mixed incontinence, frequency, incomplete voiding, and voiding difficulties were the other reported complications. SUI, also known as spontaneous urinary incontinence, is the unintentional leakage of urine during actions that raise intraabdominal pressure, such as straining, lifting, or coughing.

In women, urethral hypermobility and intrinsic sphincter weakness are the main causes of stress incontinence [26]. Buchsbaum also reported that in both community and care facility settings, elderly people frequently experience urinary incontinence and pelvic organ prolapse [26]. In cases of cystocele, urinary incontinence, particularly the SUI, is a well-recognized symptom. In comparison, Khan *et al.* observed a similar incidence of SUI (14.78%, n=69) [27], Luka *et al.* observed SUI in 15% (n=4458) of women and overactive bladder in 13% (n=4458) of their cases [28]. Bai *et al.*, however, found SUI in 63.3% (n=19) of the population [29].

The degree of the cystocele, the existence of SUI, the degree of genital prolapse, and the desire for future childbearing dictated the type of intervention. This study found that UTI, cystitis, and pyelonephritis were the most common postoperative complication. It is now standard practice to use a postoperative indwelling catheter for a limited time following prolapse surgery. On the other hand, placing an indwelling catheter increases the

chance of developing a urinary tract infection, and the rate of infection rises the longer the catheter is left in place [30].

One study by Bruce *et al.* [24] reported incontinence recurrence and voiding dysfunction after cystocele repair. Everyone who encounters women with pelvic prolapse and SUI should make an effort to offer anatomic surgical correction when necessary. We are aware that the amount of follow-up we have is little and that many continence operations will have positive outcomes right away. We need to monitor patients for a longer time to see if our defect-focused anatomic approach to incontinence and prolapse will "stand the test of time." [24]

Complications from surgical procedures are the most common. The prolapse returning and requiring additional surgery is the most frequent. After an anterior colporrhaphy, over 40% of women need a second surgery because their symptoms have returned. Other issues can result from surgery, such as postoperative bleeding, hematoma, harm to the surrounding pelvic organs, infection at the surgical site and wound dehiscence, urinary tract infection, pain during erections, urine retention, and the development of a vesicovaginal fistula [31].

The presence of a cystocele alone does not represent a risk to life. The problem is that it progresses and can result in a wide range of symptoms and illnesses if not treated properly. Women may experience voiding difficulty and urinary symptoms as a result of the bladder wall protruding into the vagina. These symptoms may be linked to kidney damage and urinary tract infections. The vaginal mucosal tissue is more prone to bleeding and infection if the prolapse extends past the vaginal entrance and causes ulcers. Additionally crucial is the psychological toll that incontinence and sexual dysfunction can have [32].

## CONCLUSION

This systematic review revealed that SUI affected the majority of cystoceles cases. The other complications included frequency, incomplete voiding, urge incontinence, mixed incontinence, and voiding difficulties. Pyelonephritis, cystitis, and UTI were the most frequent postoperative complications.

**ACKNOWLEDGMENTS :** Many thanks to Dr. Hatim T. O. Ali Assistant Professor of Obstetrics and Gynecology, College of Medicine, King Khalid University, Abha, Saudi Arabia, for his continuous help, support and encouragement to complete this work.

**CONFLICT OF INTEREST :** None

**FINANCIAL SUPPORT :** None

**ETHICS STATEMENT :** None

## REFERENCES

1. Naldini G, Fabiani B, Sturiale A, Simoncini T. Complex pelvic organ prolapse: decision-making algorithm. *Int J Colorectal Dis.* 2019;34(1):189-92.
2. Wyndaele M, Madhu C, Hashim H. Female stress urinary incontinence surgery: 'Resurgence of the Titans'. *Turk J Urol.* 2021;47(3):210-5. doi:10.5152/tud.2021.21073
3. Ghanbari Z, Ghaemi M, Shafiee A, Jelodarian P, Hosseini RS, Pouyamoghaddam S, et al. Quality of Life Following Pelvic Organ Prolapse Treatments in Women: A Systematic Review and Meta-Analysis. *J Clin Med.* 2022;11(23):7166. doi:10.3390/jcm11237166
4. Giagio S, Innocenti T, Salvioli S, Lami A, Meriggiola MC, Pillastrini P, et al. Completeness of exercise reporting among randomized controlled trials on pelvic floor muscle training for women with pelvic organ prolapse: A systematic review. *Neurourol Urodyn.* 2021;40(6):1424-32. doi:10.1002/nau.24712
5. Ruffolo AF, Lambert B, Lallemand M, Candiani M, Salvatore S, Cosson M. The Vaginal Patch Plastron Associated to the Anterior Sacrospinous Ligament Fixation for the Treatment of Advanced Anterior Vaginal Wall Prolapse. *J Clin Med.* 2022;11(22):6684. doi:10.3390/jcm11226684
6. Wallace S, Gurland B. Approaching Combined Rectal and Vaginal Prolapse. *Clin Colon Rectal Surg.* 2021;34(05):302-10. doi:10.1055/s-0041-1726351
7. Robinson D, Prodigalidad LT, Chan S, Serati M, Lozo S, Lowder J, et al. International Urogynaecology Consultation chapter 1 committee 4: patients' perception of disease burden of pelvic organ prolapse. *Int Urogynecol J.* 2022;33(2):189-210. doi:10.1007/s00192-021-04997-3

8. Durnea CM, Pergialiotis V, Duffy JM, Bergstrom L, Elfituri A, Doumouchtsis SK. CHORUS ,an International Collaboration for Harmonising Outcomes, Research and Standards in Urogynaecology and Women's Health. A systematic review of outcome and outcome-measure reporting in randomised trials evaluating surgical interventions for anterior-compartment vaginal prolapse: a call to action to develop a core outcome set. *Int Urogynecol J.* 2018;29(12):1727-45. doi:10.1007/s00192-018-3781-5
9. Shahveh M, Tajbakhsh E, Momtaz H, Ranjbar R. Antimicrobial Resistance, Biofilm Formation and Virulence Factors in *Enterococcus faecalis* Strains Isolated from Urinary Tract Infection in Kermanshah, Iran. *Arch Pharm Pract.* 2020;11(3):79-88.
10. Alsulami FT, Al-Malki YA, Al Hamad MA, Alanzi FA, Abualshamat MM, Alharbi FO, et al. Role of Family Physician in Diagnosis and Management of Urinary Tract Infection in PHC. *Arch Pharm Pract.* 2019;10(4):38-42.
11. Kalani L, Rashidi N, Mehranfard S, Bahrami H, Majidipour N, Moghaddam AS, et al. Epidemiology of the urinary stones: a 6-year retrospective study at Dezful-Iran. *Int J Pharm Phytopharmacol Res.* 2020;10(4):79-85.
12. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(1):1-0.
13. Shek KL, Rane A, Goh J, Dietz HP. Stress urinary incontinence after transobturator mesh for cystocele repair. *Int Urogynecol J.* 2009;20(4):421-5.
14. Eboue C, Marcus-Braun N, von Theobald P. Cystocele repair by transobturator four arms mesh: monocentric experience of first 123 patients. *Int Urogynecol J.* 2010;21(1):85-93.
15. Cugudda A, Terrone C, Crivellaro S, Rossetti SR. Long term results of Burch colposuspension and anterior colpoperineorrhaphy in the treatment of stress urinary incontinence and cystocele. In *Annales d'urologie* 2002 Jan 1 (Vol. 36, No. 3, pp. 176-181). Elsevier Masson.
16. Chen CH, Hsiao SM, Chang TC, Wu WY, Lin HH. Transvaginal cystocele repair using pursestring technique reinforced with custom-tailored two-armed mesh. *Urology.* 2011;78(6):1275-80.
17. Onarıminın EZ. Effect of Simultaneous Cystocele Repair On Transobturator Tape In Patients With Stress Urinary Incontinence. *J Urol Surg.* 2014;1:28-31.
18. Abdel Aziz AY, Hassan S, Morsy S, Hussein HA, Abdel Azim D, Abdel Azim MS. Outcomes of the association of asymptomatic cystocele repair in patients undergoing mid-urethral sling surgery for the correction of stress incontinence. *Afr J Urol.* 2020;26(1):1-5.
19. Park HK, Paick SH, Lho YS, Choo GY, Kim HG, Choi J. Lack of effect of concomitant stage II cystocele repair on lower urinary tract symptoms and surgical outcome after tension-free vaginal tape procedure: randomized controlled trial. *Int Urogynecol J.* 2013;24(7):1123-6.
20. Zullo MA, Ruggiero A, Plotti F, Bellati F, Basile S, Mancini N, et al. Anterior colporrhaphy plus inside-out tension-free vaginal tape for associated stress urinary incontinence and cystocele. *J Minim Invasive Gynecol.* 2008;15(4):446-51.
21. Okui N, Okui M, Horie S. Improvements in overactive bladder syndrome after polypropylene mesh surgery for cystocele. *Aust N Z J Obstet Gynaecol.* 2009;49(2):226-31.
22. Shrestha S. Urinary complication following cystocele repair in pelvic organ prolapse. *Nepal J Obstet Gynaecol.* 2016;11(1):20-3.
23. Adile B, Gugliotta G, Adile G, Verso LL, Sommatino F, Scurria S. Monoprosthesis for simultaneous correction of stress urinary incontinence and cystocele: a 3-year follow-up. *Urol J.* 2012;79(19\_suppl):134-7.
24. Bruce RG, El-Galley RE, Galloway NT. Paravaginal defect repair in the treatment of female stress urinary incontinence and cystocele. *Urology.* 1999;54(4):647-51.
25. Gilleran JP, Lemack GE, Zimmern PE. Reduction of moderate-to-large cystocele during urodynamic evaluation using a vaginal gauze pack: 8-year experience. *BJU Int.* 2006;97(2):292-5.
26. Jaffar A, Mohd-Sidik S, Nien FC, Fu GQ, Talib NH. Urinary incontinence and its association with pelvic floor muscle exercise among pregnant women attending a primary care clinic in Selangor, Malaysia. *PLoS One.* 2020;15(7):e0236140. doi:10.1371/journal.pone.0236140
27. Sori DA, Bretones S, Mellier G, de Rochambeau B. Prevalence and surgical outcomes of stage 3 and 4 pelvic organs prolapse in Jimma university medical center, south west Ethiopia. *BMC Women's Health.* 2022;22(1):1-0. doi:10.1186/s12905-022-01992-8

28. Zafarnia M, Kennes LN, Stickeler E, Hoff J, Najjari L. Evaluation of urinary continence status and its influence on quality of life after gynecological treatment of female pelvic malignancies at an oncological center. *BMC Women's Health*. 2022;22(1):1-0. doi:10.1186/s12905-022-01999-1
29. Harvey MA, Chih HJ, Geoffrion R, Amir B, Bhide A, Miotla P, et al. International Urogynecology Consultation Chapter 1 Committee 5: relationship of pelvic organ prolapse to associated pelvic floor dysfunction symptoms: lower urinary tract, bowel, sexual dysfunction and abdominopelvic pain. *Int Urogynecol J*. 2021;32(10):2575-94. doi:10.1007/s00192-021-04941-5
30. Xie N, Hu Z, Ye Z, Xu Q, Chen J, Lin Y. A systematic review comparing early with late removal of indwelling urinary catheters after pelvic organ prolapse surgery. *Int Urogynecol J*. 2021;32(6):1361-72. doi:10.1007/s00192-020-04522-y
31. Dietz HP, Hankins KJ, Wong V. The natural history of cystocele recurrence. *Int Urogynecol J*. 2014;25(8):1053-7.
32. Makajeva J, Watters C, Safioleas P. Cystocele. InStatPearls [Internet] 2021 Sep 18. StatPearls Publishing.