



Original Article

ISSN : 2277-3657
CODEN(USA) : IJPRPM

Effect of Smokeless Tobacco (Mishri) Application on Periodontal Health and Correlation to the Duration of Use

**Girish Suragimath^{1*}, Jay Narendra Patel¹, Tanvi Sandeep Mhatre¹, Ashwinirani SR²,
Dhirajkumar Arun Mane³**

¹Department of Periodontology, School of Dental Sciences, Krishna Institute of Medical Sciences Deemed University, Karad, Maharashtra, India.

²Department of Oral Medicine and Radiology, School of Dental Sciences, Krishna Institute of Medical Sciences Deemed University, Karad, Maharashtra, India.

³Directorate of Research Department, Krishna Institute of Medical Sciences Deemed University, Karad, Maharashtra, India.

*Email: drgirishsuragimath@gmail.com

ABSTRACT

Mishri is a form of smokeless tobacco used as a dentifrice in rural areas of central India. Mishri is applied to teeth and gums to clean the oral cavity. Mishri use has ill effects on oral health and periodontal tissues. The present study assessed the ill effect of mishri application periodontal tissues, and its correlation to the duration of use. A total of 300 subjects who used mishri as a dentifrice were considered. The enrolled subjects were categorized into three subgroups depending on the years of mishri usage. Age, gender, education level, socioeconomic status (SES), and body mass index (BMI), were recorded. Periodontal clinical parameters i.e. plaque index, gingival index, clinical attachment loss, and probing pocket depth were assessed. The data were statistically correlated with the duration of mishri usage with the Chi-square test and Analysis of variance. Out of the 300 subjects, 162 (54%) were males and 138 (46%) were females. SES and BMI had no association with mishri use but education level played a significant role in the avoidance of mishri use ($P = 0.041$). All the periodontal clinical parameters deteriorated with the duration of mishri use ($P = 0.0001$). It was concluded that mishri use was common in both genders. There was a deterioration of periodontal health and the progressive destruction of periodontal tissues with years of mishri use. The Dentist and community health workers should educate about the ill effects of tobacco use and mishri, as a dentifrice among the rural population.

Key words: *Clinical attachment loss, Gingival health, Mishri, Periodontal disease, Smokeless tobacco*

INTRODUCTION

The use of smokeless tobacco in different forms is rampant throughout India. Betel quid chewing and smokeless tobacco use is a socially accepted practice and has become part of cultural and religious customs. Smokeless tobacco is available throughout India in different forms like chewing, snuffing, application to teeth and gums, Paan, Khaini, Zarda, Mishri, Gutkha, Toombak etc. Oral smokeless tobacco products are placed in the mouth, cheek or lip and sucked or chewed [1]. Smokeless tobacco (SLT) is defined as a product that contains tobacco, is not smoked or burned at the time of use, and is commonly consumed orally or nasally [2].

India harbors about 70% of the world's SLT users, which has the potential to cause premalignant, malignant cancers and poor reproductive outcomes [3]. Mishri is a form of SLT that is used as a dentifrice to clean teeth in the central part of India. Mishri is mostly a homemade preparation, by roasting tobacco leaves on a hot metal plate

till it becomes black uniformly, then powdered and packed. Mishri is also commercially available locally in different trade names [4]. Various studies have proved the ill effects of SLT on dental and periodontal tissues like gingival recession, loss of attachment, discolouration of teeth, tooth abrasion, halitosis and dental implant failures [5-8]. There is a need to study the ill effects of mishri application on the periodontal tissues. With this background, the present study was designed to assess the ill effect of mishri use on periodontal tissues, and its correlation to the duration of use.

MATERIALS AND METHODS

This cross-sectional study was conducted at the department of Periodontology, School of Dental Sciences, Karad, Maharashtra, India. A total of 300 subjects who used mishri as a dentifrice to clean their teeth were enrolled in the study by simple random purposive sampling technique. Institutional ethical clearance was obtained from Krishna Institute of Medical Sciences Deemed to be University (KIMSDU) before commencing the study (Letter No: KIMSDU/IEC/03/2018). The objective of the study was explained to all the enrolled subjects, and written informed consent was obtained from each of them. Demographic data (education level, occupation and socioeconomic status) were recorded in a structured proforma.

Subjects aged 19 -72 years with the presence of at least 20 permanent teeth, were considered for the study. Pregnant and lactating women and subjects suffering from any systemic diseases or using any other form of tobacco, and also those who had undergone periodontal therapy within the last six months were excluded from the study.

The enrolled subjects were categorized into three subgroups depending on the years of mishri usage. Group 1: Subject using mishri for less than 5 years, Group 2: Subject using mishri from 5 to 10 years, and Group 3: Subjects using mishri for more than 10 years.

A detailed history of mishri usage from each participant was recorded, which included, the number of mishri applications per day and years of usage. The socioeconomic status (SES) of the participants was assessed by the Modified Kuppaswamy SES scale 2021 and divided into five categories 1 - Lower, 2 - Upper Lower, 3 - Lower Middle, 4 - Upper Middle, and 5 – Upper [9, 10]. The body mass index of the participants was calculated using the formula $BMI = kg/m^2$ and graded as 1 - underweight $BMI < 18.5 kg/m^2$, 2 - normal weight $BMI = 18.5$ to 23, 3 - overweight $BMI = 23$ to 25 and 4 - obesity $BMI > 25$ [11]. The education level of the participants was enquired and divided into seven grades 1 - Illiterate, 2 – Primary school certificate, 3 – Middle school certificate, 4 – High school certificate, 5 – Post high school diploma, 6 – Graduate or postgraduate and 7 – Professional degree [11]. Clinical examination of the periodontal tissues was carried out by a single trained calibrated operator. All the subjects enrolled in the study were subjected to periodontal analysis which included the Gingival index, and Plaque Index [12, 13]. Probing pocket depth (PPD) and Clinical attachment level (CAL) were recorded to the nearest millimetre with a periodontal probe (Williams periodontal probe Hu-Friedy, Rotterdam, Netherland) at six sites (mesiobuccal, buccal, distobuccal, mesiolingual, lingual and distolingual) of all teeth excluding the third molars [14].

The data obtained were compiled and entered into Microsoft Excel 2010. The Chi-square test and analysis of variance (ANOVA) were used to find the association and significance. Statistical analyses were performed using statistical software IBM SPSS Statistics for Windows, version 20.0 (IBM Corp., Armonk, N.Y., USA), and a P value < 0.05 was considered statistically significant.

RESULTS AND DISCUSSION

A Total of 300 subjects who used mishri as a dentifrice were included in the study, consisting of 162 males (54%) and 138 females (46%). The mean age of the participants was 40.6 ± 4.2 years (**Table 1**).

Table 1. Gender and age of the study subjects.

Gender	N [%]	Age Mean \pm SD
Male	162 (54%)	40.62 \pm 3.84
Female	138 (46%)	40.57 \pm 4.82
Total	300 (100%)	40.60 \pm 4.20

The participants were categorized into three groups on the duration of the mishri usage in years: group 1: under five years, group 2: six to ten years, and group 3: more than ten years. There was no association found between gender and duration of mishri usage P-value: 0.99 (**Table 2**).

Table 2. Study subjects categorized according to gender and duration mishri usage

Groups	Duration of Mishri Use	Males N (%)	Females N (%)	Total N (%)	χ^2 Value	P-value
Group 1	under 5 years	74 (45.68%)	67(48.55%)	141 (47.00%)	0.25	0.99
Group 2	6-10 years	56 (34.57%)	45(32.61%)	101 (33.67%)		
Group 3	more than 10years	32 (19.75%)	26 (18.84%)	58 (19.33%)		
Total		162 (100%)	138 (100%)	300 (100%)		

The education level, socioeconomic status, and Body Mass Index of the participants were recorded and the Chi-square test was used to find the association between the mishri uses. The education level had a significant association with mishri use P-value: 0.0413. The socioeconomic status P-value: 0.42 and BMI of the subjects had no association with mishri use P-value: 0.072 (**Table 3**).

Table 3. Education level, socioeconomic status, and body mass index of the participants.

Groups	Group 1 N (%)	Group 2 N (%)	Group 3 N (%)	χ^2 Value	p-value
Education level					
1 – Illiterate	01 (0.33)	02 (0.67)	01 (0.33)	21.68	0.0413*
2 – Primary school certificate	13 (4.33)	17 (5.67)	18 (6.00)		
3 – Middle school certificate	12 (4.00)	15 (5.00)	18 (6.00)		
4 – High school certificate	41 (13.67)	28 (9.33)	35 (11.67)		
5 – Post high school diploma	29 (9.67)	21 (7.00)	19 (6.33)		
6 – Graduate or postgraduate	18 (6.00)	04 (1.33)	02 (0.67)		
7 – Professional degree	03 (1.00)	02 (0.67)	01 (0.33)		
Socioeconomic status					
1 – Lower	51 (17.00)	38 (12.67)	36 (12.00)	8.09	0.42
2 - Upper lower	22 (7.33)	16 (5.33)	24 (8.00)		
3 - Lower middle	11 (3.67)	16 (5.33)	18 (6.00)		
4 - Upper middle	10 (3.33)	13 (4.33)	10 (3.33)		
5 -Upper	09 (3.00)	11 (3.67)	15 (5.00)		
BMI					
1 - Underweight	16 (5.33)	18 (6.00)	18 (6.00)	11.55	0.072
2 - Normal	71 (23.67)	62 (20.67)	41 (13.67)		
3 - Overweight	17 (5.67)	14 (4.67)	10 (3.33)		
4 – Obese	10 (3.33)	25 (8.33)	08 (2.67)		

*Significant p<0.05

The periodontal status of the study subjects was assessed using the following clinical parameters plaque index, gingival index, clinical attachment loss, and probing pocket depth. ANOVA was used to correlate the periodontal status with the duration of mishri use. All the clinical parameters had a statistically significant correlation with the duration of mishri use with a P-value of <0.0001 (**Table 4**).

Table 4. Periodontal clinical parameters of the study subjects.

Groups	Group 1 Mean±SD	Group 2 Mean±SD	Group 3 Mean±SD	ANOVA F-value	p-value
Plaque index	1.03±0.21	1.50±0.45	1.74±0.54	218.12	<0.0001*
Gingival index	1.27±0.34	1.40±0.09	1.52±0.38	52.48	<0.0001*

Clinical attachment loss (mm)	2.45±0.74	3.23±0.46	4.27±0.18	947.9	<0.0001*
Probing pocket depth (mm)	2.08±0.45	2.78±0.46	3.95±0.08	1910.5	<0.0001*

*Significant p<0.05

The negative effect smoked form of tobacco on general and periodontal health has been extensively researched, documented and proven. Cigarette smoke induces oxidative stress and is involved excessive inflammatory response of human pulmonary artery smooth muscle cells, and sexual dysfunction through testicular damage [15-17]. The literature regarding the ill effects of a smokeless form of tobacco on periodontal tissues is sparse. The present cross-sectional study was designed to assess the ill effect of smokeless tobacco (mishri) use on periodontal tissues and its correlation to the duration of use.

The use of mishri as a dentifrice was mostly observed in the rural population with a mean age of 40.6 ± 4.2 years. The results of our study are similar to the previous studies by Katturi *et al.*, who found that 20 to 40 years of age subjects used tobacco in a smoked and smokeless form. The age difference of the higher age group in our study may be due to, we have considered subjects only using mishri as a dentifrice for our study, and excluded other forms of SLT users [18].

The use of Mishri as a dentifrice was almost equal in both gender, males (54%) and females (46%). The results were similar to the study conducted in Bangladesh which found the smokeless form of tobacco use was equally prevalent in males and females [19]. Shah *et al.*, the study concluded that SLT use among Indian females is equally prevalent as the male population [20].

The use of Mishri as a dentifrice for years was observed in many individuals. There was no correlation seen statistically (P-value: 0.99) according to the duration of use of mishri. It was evident that the study subjects used mishri for more than ten years without the knowledge and awareness of its ill effects on the periodontium and oral cavity.

The education level of the study subjects played a key role in the use of mishri in the study population. Subjects with higher education i.e. post high school diploma, graduate and professional degree refrained from using mishri as a dentifrice (P-value: 0.0413). The results of our studies in accordance with previous literature, that education level played an important role in smokeless tobacco use [21, 22]. A cohort study on the Sweden population, concluded that tobacco product users were less educated compared to non-users [23].

The association between socioeconomic status and BMI with the mishri use was not statistically significant (P-value: 0.42). The data from our study show that even upper and upper-middle-class subjects used mishri as a dentifrice in the Maharashtra region. The results of our study are in accordance with Jodalli PS, Panchmal GS Study on the Indian population, who concluded that socioeconomic status did not correlate with tobacco use [24]. The BMI had no association with mishri use in our study (P-value: 0.072), which is in accordance with the previous research conducted on the Indian population [25]. The present study subjects were mostly rural populations from farming backgrounds who were physically active and were not obese.

All the periodontal clinical parameters in our study population increased with the years of mishri use denoting that periodontal disease status deteriorated with years of mishri use, which is in accordance with previous research on the use of SLT [26, 27]. The plaque levels and gingival inflammation increased in our study subjects with the duration of mishri use. The results are similar to previous research by Offenbacher S and Weathers DR study [28, 29]. The CAL and PPD increased with the duration of the mishri use in our study which was statistically significant. The results of our study are similar to previous studies, where they found progressive destruction of periodontal tissues in SLT users [30-32].

Quitting the use of tobacco is difficult, and not an easy task for a regular users, due to its addictive properties [33]. Tobacco counselling with the guidance from physician, dentist, health workers and family members can help the tobacco user to quit the tobacco habit. There is clear and urgent need by all the countries to reduce the prevalence of tobacco use, both in smoked and smokeless forms, by passing strong evidence based policies, which will benefit the general population at large [34].

CONCLUSION

The results of our study denote that the use of mishri as a dentifrice was common among middle age and older individuals in both male and female gender. The level of education played a key role in the avoidance of mishri use as a dentifrice. There was a deterioration of periodontal health and the progressive destruction of periodontal tissues with years of mishri use.

The Dentist and community health workers should educate the general population about the ill effects of tobacco use and mishri as a dentifrice. Tobacco cessation and health education promotions should be carried out at the rural level for benefit of the general population.

ACKNOWLEDGMENTS : None

CONFLICT OF INTEREST : None

FINANCIAL SUPPORT : None

ETHICS STATEMENT : The Ethical clearance for the study was obtained from the Institutional Ethical Committee (IEC) of KIMSDU, Karad, (Letter No: KIMSDU/IEC/03/2018).

REFERENCES

1. Dobe M, Sinha DN, Rahman K. Smokeless tobacco use and its implications in WHO South East Asia Region. *Indian J Public Health*. 2006;50(2):70-5.
2. Singh S, Jain P, Singh PK, Reddy KS, Bhargava B. White paper on smokeless tobacco & women's health in India. *Indian J Med Res*. 2020;151(6):513-21. doi:10.4103/ijmr.IJMR_537_20
3. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419-31. doi:10.1046/j.1440-1843.2003.00507.x
4. Pratinidhi A, Gandham S, Shrotri A, Patil A, Pardeshi S. Use of 'Mishri' A Smokeless form of Tobacco During Pregnancy and its Perinatal Outcome. *Indian J Community Med*. 2010;35(1):14-8. doi:10.4103/0970-0218.62547
5. Montén U, Wennström JL, Ramberg P. Periodontal conditions in male adolescents using smokeless tobacco (moist snuff). *J Clin Periodontol*. 2006;33(12):863-8. doi:10.1111/j.1600-051X.2006.01005.x
6. Frithiof L, Anneroth G, Lasson U, Sederholm C. The snuff-induced lesion. A clinical and morphological study of a Swedish material. *Acta Odontol Scand*. 1983;41(1):53-64. doi:10.3109/00016358309162303
7. Verma SK, Kumar BD, Singh S, Kumari P, Agarwal A, Singh TK, et al. Effect of gutkha chewing on periodontal health and oral hygiene of peoples in Delhi NCR region of North India: A cross-sectional multicentered study. *J Family Med Prim Care*. 2019;8(2):564-7. doi:10.4103/jfmpe.jfmpe_439_18
8. Mittal S, Dani N, Abullais SS, Al-Qahtani NA, Shah K. Effect of Smoking and Tobacco Chewing on Periodontal Disease and Non-Surgical Treatment Outcome: A Clinical and Biochemical Study. *J Int Acad Periodontol*. 2017;20(1):12-8.
9. Saleem SM, Jan SS. Modified Kuppuswamy socioeconomic scale updated for the year 2021. *Indian J ForensicCommunity Med*. 2021;8(1):1-3.
10. Wani RT. Socioeconomic status scales-modified Kuppuswamy and Udai Pareekh's scale updated for 2019. *J Family Med Prim Care*. 2019;8(6):1846-9. doi:10.4103/jfmpe.jfmpe_288_19
11. Weir CB, Jan A. BMI classification percentile and cut off points. 2021. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022.
12. Løe H, Silness J. Periodontal disease in pregnancy I. Prevalence and severity. *Acta Odontol Scand*. 1963;21(6):533-51. doi:10.3109/00016356309011240
13. Silness J, Loe H. Periodontal disease in pregnancy. II. correlation between oral hygiene and periodontal condition. *Acta Odontol Scand*. 1964;22:121-35. doi:10.3109/00016356408993968
14. Carranza FA, Newman MG, Takkie HH, Klokkevold PR. *Clinical Periodontology*. 7th ed. Vol. 2. 2013. pp. 487-491.
15. Wang J, Wang L, Chen X, Liang ML, Wei DH, Cao J, et al. Cigarette smoke extract stimulates human pulmonary artery smooth muscle cell proliferation: Role of inflammation and oxidative stress. *Iran J Basic Med Sci*. 2022;25(6):755-61. doi:10.22038/IJBMS.2022.64170.14133
16. Nazmara Z, Ebrahimi B, Makhdoumi P, Noori L, Mahdavi SA, Hassanzadeh G. Effects of illicit drugs on structural and functional impairment of testis, endocrinal disorders, and molecular alterations of the semen. *Iran J Basic Med Sci*. 2021;24(7):856-67. doi:10.22038/ijbms.2021.53326.12002
17. Alalwani AD. Protective effect of grape juice on testicular and epididymis damage in rats exposed to cigarette smoke. *Int J Pharm Phytopharmacol Res*. 2020;10(3):185-93.

18. Katuri KK, Alluri JK, Chintagunta C, Tadiboina N, Borugadda R, Loya M, et al. Assessment of periodontal health status in smokers and smokeless tobacco users: a cross-sectional study. *J Clin Diagnostic Res.* 2016;10(10):ZC143-6.
19. Flora MS, Mascie-Taylor CG, Rahman M. Gender and locality differences in tobacco prevalence among adult Bangladeshis. *Tob Control.* 2009;18(6):445-50. doi:10.1136/tc.2008.028142
20. Shah S, Dave B, Shah R, Mehta TR, Dave R. Socioeconomic and cultural impact of tobacco in India. *J Family Med Prim Care.* 2018;7(6):1173-6. doi:10.4103/jfmpc.jfmpc_36_18
21. Sharma D, Goel S, Lal P. Education differential in relation to tobacco use and its predictors across different regions of India. *Indian J Cancer.* 2017;54(3):584-8. doi:10.4103/ijc.IJC_345_17
22. Thakur JS, Paika R. Determinants of smokeless tobacco use in India. *Indian J Med Res.* 2018;148(1):41-5. doi:10.4103/ijmr.IJMR_27_18
23. Julkunen-Iivari A, Heikkinen AM, Räisänen IT, Ruokonen H, Meurman JH, Toppila-Salmi S, et al. Tobacco Products, Periodontal Health and Education Level: Cohort Study from Sweden. *Dent J (Basel).* 2020;8(3):90. doi:10.3390/dj8030090
24. Jodalli PS, Panchmal GS. Socioeconomic Correlates and Key Aspects of Tobacco Surveillance Using Global Adult Tobacco Survey Among College Students of Mangaluru, South India. *Cureus.* 2019;11(2):e4115. doi:10.7759/cureus.4115
25. Pednekar MS, Gupta PC, Shukla HC, Hebert JR. Association between tobacco use and body mass index in urban Indian population: implications for public health in India. *BMC Public Health.* 2006;6:70. doi:10.1186/1471-2458-6-70
26. Anand PS, Mishra S, Nagle D, Kamath NP, Kamath KP, Anil S. Patterns of Periodontal Destruction among Smokeless Tobacco Users in a Central Indian Population. *Healthcare (Basel).* 2021;9(6):744. doi:10.3390/healthcare9060744
27. Goel K, Sharma S, Baral DD, Agrawal SK. Current status of periodontitis and its association with tobacco use amongst adult population of Sunsari district, in Nepal. *BMC Oral Health.* 2021;21(1):66. doi:10.1186/s12903-021-01416-3
28. Offenbacher S, Weathers DR. Effects of smokeless tobacco on the periodontal, mucosal and caries status of adolescent males. *J Oral Pathol.* 1985;14(2):169-81. doi:10.1111/j.1600-0714.1985.tb00480.x
29. Mahapatra S, Chaly PE, Mohapatra SC, Madhumitha M. Influence of tobacco chewing on oral health: A hospital-based cross-sectional study in Odisha. *Indian J Public Health.* 2018;62:282-6.
30. Yaragani A, Sushuma K, Guduri V, Thirumalasetty SSMK, Vishnubhotla G, Kandikatla P, et al. The influence of tobacco consumption on periodontal health: A stratified analysis based on type of tobacco use. *J Family Med Prim Care.* 2020;9(4):2061-6. doi:10.4103/jfmpc.jfmpc_1071_19
31. Katuri KK, Alluri JK, Chintagunta C, Tadiboina N, Borugadda R, Loya M, et al. Assessment of Periodontal Health Status in Smokers and Smokeless Tobacco Users: A Cross-Sectional Study. *J Clin Diagn Res.* 2016;10(10):ZC143-6. doi:10.7860/JCDR/2016/22160.8700
32. Ahad A, Bey A, Khan S, Ahmad MS. Periodontal status associated with dual habits of smoking and smokeless tobacco use: A cross-sectional study in young adults. *J Adv Periodontol Implant Dent.* 2021;13(2):69-75. doi:10.34172/japid.2021.010
33. Bozkurt N, Altıntas F, Bozkurt AI, Turgut G, Turgut S. Effect of MDR C3435T polymorphism on Varenicline treatment in quit smoking. *Braz J Pharm Sci.* 2019;550:e18186.
34. GBD 2019 Tobacco Collaborators. Spatial, temporal, and demographic patterns in prevalence of smoking tobacco use and attributable disease burden in 204 countries and territories, 1990-2019: a systematic analysis from the Global Burden of Disease Study 2019. *Lancet.* 2021;397(10292):2337-60. doi:10.1016/S0140-6736(21)01169-7