



Research Article

ISSN : 2277-3657  
CODEN(USA) : IJPRPM

## ***Effectiveness of Aqueous Extract of Fenugreek Seeds and Flaxseed on Polycystic Ovarian Syndrome in Female Rats***

**Nadia Nour Osman<sup>1,2</sup>, Shyma Ali Alsahfi<sup>1\*</sup>, Fawzia Alshubaily<sup>1</sup>**

<sup>1</sup>Department of Biochemistry, Faculty of Science, King Abdulaziz University, P. O. 42801, Jeddah - 21551, Saudi Arabia.

<sup>2</sup>Department of Food Irradiation Research, National Centre for Radiation Research and Technology (NCRRT), Cairo, Egypt.

\*Email: [shyma.a.sahafi@gmail.com](mailto:shyma.a.sahafi@gmail.com)

---

### **ABSTRACT**

Polycystic ovarian syndrome (PCOS) is a disease impacting adult female in which its main features are hyperandrogenism, insulin resistance (IR), hyperinsulinemia, which enhances the probability of getting type 2 diabetes and cardiovascular disease. This study aimed to estimate the antidiabetic effect of fenugreek and the anti-androgenic effect of flaxseed and use the combination of both to treat polycystic ovarian syndrome in female rats. Forty Wistar rats were divided into five groups each contained eight rats, Group I: normal control, the rats in Group II-V received Estradiol valerate (4mg/kg in 0.4ml Sesame oil i.p.). Then, Group II served as PCOS control. The rats in Group-III were given fenugreek aqueous extract (100 mg/kg), Group IV received flaxseed aqueous extract (300 ml/kg) and Group V administered a combination of both extracts, for 60 consecutive days after PCOS conformation, extracts were given orally. The results showed that the Estradiol Valerate induced a dignified increase in glucose, insulin, insulin resistance, lipid profile, aspartate aminotransferase (AST), alanine aminotransferase (ALT), gamma-glutamyltranspeptidase (GGT), and body weight in comparison to control. Moreover, an outstanding rise in serum LH, FSH, testosterone and a decrease in progesterone level, with no change in estradiol was detected. Furthermore, a significant rise in lipid peroxidation accompanied by an insignificant increase in nitric oxide, a significant decline in superoxide dismutase, catalase, and glutathione reductase activities in ovarian homogenates were also noted. Supplementation of fenugreek and flaxseed significantly attenuated these parameters. The results showed the beneficial effect of fenugreek and flaxseed in improving the impairment of IR and hyperandrogenism with corresponding disrupted parameters in polycystic ovarian syndrome.

**Key words:** *Hyperandrogenism, hyperinsulinemia, Insulin Resistance, Estradiol Valerate.*

---

### **INTRODUCTION**

Polycystic ovarian syndrome (PCOS) form the greatest endocrinal disorders that influence women after reaching their procreation period age, [1] affecting around one of fifteen women widespread, distinguished by raised insulin and androgen levels, [2] which at latest leads to acne and hirsutism. It is also associated with insulin resistance (IR), obesity,[3] anovulation, and infertility on the prolonged incidence of cysts[4]. PCOS has been shown to be related to metabolic syndrome and its abnormalities [5, 6]. The major used diagnosis criteria of PCOS is Rotterdam criteria [7]. Clinical diagnostic criteria of PCOS is suggested to be the presence of hirsutism, hyperandrogenism, irregular menses, [8] metabolic syndrome characteristics, anovulation, and the polycystic ovaries [9]. PCOS etiology is not well recognized yet and trial animal models may assist to understand the pathophysiology of PCOS [10]. PCOS is correlated with hyperlipidemia and glucose intolerance in having a risk factor of type 2 diabetes, hypertension, cardiovascular disorder, and cancer [11]. Pharmaceutical medicines used for decades to treat PCOS include insulin sensitizers such as metformin and pioglitazone for hyperinsulinemia and hyperandrogenism, [12] oral contraceptives for hirsutism and menstrual irregularities, Spironolactone and finasteride for increased androgen, and clomiphene for infertility [13]. Complementary and alternative medicine

(CAM) is globally improving and preferred by individuals [14, 15]. There is demanding use of CAM therapies in an alternative to drugs for chronic diseases [16]. Fenugreek ("*Trigonella foenum-graecum L.*") is a leguminous, annual herb growing in India and North Africa [17]. It showed an anticancer effect against human hepatoma, [18] antidiabetic properties, [19] hypocholesterolemic effect, [20] and anti-inflammatory activity against arthritis in rats [21]. Flaxseed or linseed (*Linum usitatissimum L.*), it belongs to Linaceae family [22]. Flaxseed is loaded with bioactive components, which are efficient in the protection of some long-term illnesses like numerous types of cancer, cardiovascular diseases and cerebrovascular stroke, [23] and diabetes [24]. It is also rich in lignans, which showed to decline the level of androgen in males with prostate cancer [25]. This study was planned to assess the antidiabetic effect of fenugreek, the antiandrogenic effect of the flaxseed, and the effect of their combination in the treatment of polycystic ovary syndrome in female rats.

## **MATERIALS AND METHODS:**

### **Experimental Animals**

In this experiment, 40 female Wistar rats with the weight range between 170-200g were obtained from the Animal House of King Fahd Research Center, Jeddah, Saudi Arabia. The rats were kept at normal temperature and standard light cycle for one week before the experiment began to adapt with adequate food and water. The Animal Ethics Committee (Reference No 580-17) approved the experimental animal procedure in accordance with the guidelines for proper care and use of laboratory experimental animals of King Abdulaziz University.

### **Plant Material and Preparation**

Fenugreek and flaxseed were purchased from a local traditional market in Jeddah, Saudi Arabia. 25 g of fenugreek seed was drenched in 100 ml of hot distilled water for 30 minutes, then filtered into a conical flask and the extract was collected and preserved for further usage [26]. 20 g of flaxseed [25] was soaked in 100 ml of hot distilled water and the same preparation of fenugreek was used.

### **Experimental Design**

Rats were divided into five groups, each group with 8 rats. Group I was served as the normal control group and was given only distilled water. Group II-V were intramuscularly injected with 4 mg/rat (0.4 ml) of estradiol valerate dissolved in sesame oil once to generate the PCOS model. Vaginal smears were regularly taken and monitored under the microscope to ensure PCOS formation by watching the dominant cell type during their estrous cycle. Group II was served as the positive control PCOS-induced group and was given distilled water orally during the whole experiment period. Group III was addressed orally with 100 mg/kg of fenugreek aqueous extract daily. Group IV was daily given 300 mg/kg of flaxseed aqueous extract. Group V was daily administered with both 100mg/kg of fenugreek extract and 300 mg/kg of flaxseed extract. During the experiment, the body weight was recorded every week. After 60 days of treatment, blood samples were collected after euthanasia of rats with diethyl ether. Ovaries were removed and weighed. Serum was separated using a centrifuge and used for estimation of blood glucose, insulin, lipid profile : cholesterol and triglycerides levels, FSH, LH, estradiol, progesterone, and testosterone. IR was calculated by the homeostatic model assessment (HOMA) formula. AST, ALT, GGT were also estimated to study the liver function. Oxidative stress was as well studied in ovary tissues by estimating the activity of catalase, SOD, glutathione reductase, lipid peroxidation, and nitric oxide using a special kit purchased from Elabscience Company for each antioxidant.


### **Statistical Analysis**

The data were analyzed using Megastat 10.0 add-in of Microsoft Excel program (2010). They were demonstrated as "mean  $\pm$  standard deviation (SD)" by using One-way analysis of variance (ANOVA). Values higher than or equal to 0.05 were considered no significant difference and values less than 0.05 were counted as a significant difference.

## **RESULTS:**

### **Vaginal Smear and Estrous Cycle**


An irregular estrous cycle was spotted in EV-induced rats with respect to control with a predominance of cornified cells type indicating the constancy of the estrous cycle in most rats with PCOS (Fig 1).



**Figure 1:** Vaginal smears of rats' estrous cycle showed different estrous phases with their presented type of cells (A) pro-estrous stage filled with E-epithelial cells, (B) estrous phase identified with c-cornified cells, (C) met-estrous phase with an equal amount of each cell type, and (D) diestrous stage mainly filled with L-leukocytes.

**Body and Ovarian Weight**

EV-induced rats revealed a significant increase in ovarian and body weight with respect to the control group with significant improvement after fenugreek and flaxseed supplementation (Fig 2).



**Figure 1:** The Effect of EV and treatment of fenugreek and/or flaxseed on body weight of induced rats

**Glucose, Insulin, and IR Status in Control, EV-induced and Treated Groups**

Control group had normal levels of glucose, insulin, and IR while they were significantly higher in EV-induced rats. Administration of fenugreek and flaxseed showed a significant reduction in those parameters (Table 1).

**Table 1:** The effect of fenugreek and/or flaxseed on glucose, Insulin, and Insulin Resistance in EV- induced groups


Groups	Glucose (mg/dL)	Insulin (µU/ml)	HOMA - IR
Control	80.03 ± 9.10	6.30 ± 0.59	1.25 ± 0.17
PCOS	122.56 ± 4.58	21.57 ± 2.31	6.53 ± 0.87

	*** ^^^	*** ^^^	*** ^^^
<b>PCOS + Fenugreek</b>	106.63 ± 6.82 *** ### ^^^	13.63 ± 1.79 *** ### ^^^	3.59 ± 0.54 *** ### ^^^
<b>PCOS + Flaxseed</b>	112.50 ± 6.72 *** ### ^^^	18.06 ± 2.08 *** ### ^^^	5.03 ± 0.78 *** ### ^^^
<b>PCOS + Fenugreek + Flaxseed</b>	89.75 ± 8 ### **	9.27 ± 0.97 ### ***	2.04 ± 0.30 ### **

The values are the mean ± S.D. of parameters measured  
 P < 0.01 \*\*, P < 0.001 \*\*\* significantly different from control group.  
 P < 0.001 ### significantly different from PCOS group.  
 P < 0.001 ^^ significantly different from PCOS treated with Fenugreek + Flaxseed group.

**Lipid Profile in EV- induced and Treated Groups:**


Levels of cholesterol, TG, and LDL were significantly higher in PCOS group while HDL was reduced in comparison to control. These disturbed values were attenuated after supplementation of fenugreek and flaxseed (Fig 3).



**Figure 2:**Effect of fenugreek and/or flaxseed on lipid profile parameters level in EV induced PCOS rat

**Reproductive Hormones in EV-induced and Treated Rats**

A significant increase was observed in LH, FSH, and testosterone after EV induction with a corresponding decrease in progesterone in contrast to control. On the other hand, estradiol was the same in all groups with no change. However, after administration of fenugreek and flaxseed, all FSH, LH, testosterone, and progesterone were significantly restored back to normal (Fig 4).



**Figure 3:** Effect of EV, fenugreek and/or flaxseed on the concentration of reproductive hormone parameters

**Liver Enzymes in Control, PCOS, and Treated Groups**

ALT, AST, and GGT showed a significant rise in EV-induced rats compared to those in the control, followed by a significant decrease in these liver enzymes after fenugreek and flaxseed supplementation (Fig 5)




Figure 4: Effect of EV, fenugreek and/or flaxseed on liver enzymes

**Oxidative Stress and Antioxidants in EV-induced and treated groups**

Administration of EV to female rats provokes oxidative stress in ovaries, manifested by a significant concomitant increase in the levels of LPO, an insignificant increase in NO, a significant decrease in the activities of the antioxidants enzymes, SOD, CAT, and GR in ovarian tissues. On the other hand, in PCOS rats treated with fenugreek and/or flaxseed, a significant increase in the reduced antioxidants along with a significant decrease in LPO and NO was observed compared to the corresponding values in untreated PCOS rats (Fig 6 and 7).




Figure 5: Effect of antioxidants in EV-induced and treated rats




Figure 6: Effect of oxidants in EV-induced and treated rats

**DISCUSSION:**

The current study focused on the hypoglycemic effect of fenugreek and its influence on IR reduction, as well as the anti-androgenic effect of flaxseed concentrates in PCOS-induced rats in which the symptoms were the main treatment directions in this syndrome. The results showed a significant increase in body and ovarian weights of EV treated rats. Ghafurniyan *et al.* [27], Karimzadeh *et al.* [28], Jashni & Jahromi *et al.* [29], and Patel & Shah [12] concluded that the increase in ovaries and body weights attributed to the formation of cystic follicles in the ovaries of PCOS rats and increase in body fat. The treatment of PCOS rats with fenugreek and flaxseed showed improvement and a significant decrease in ovarian and body weights. Swaroop *et al.* [30] revealed a reduction in ovarian volume in PCOS subjects after administration of fenugreek seed extract, almost half of them showed reduced cyst size while 36% showed fully cyst degeneration. Muraki *et al.* [31] showed that fenugreek is effective against obesity and it decreases body fats. Moreover, Moradi *et al.* [32] pointed to the existence of fenugreek active components such as flavonoids, steroids, saponins, and alkaloids. These constituents have mainly antioxidant effect [33, 34]. Moreover, these active contents probably participate in its effect on body and ovarian weight. Flaxseed treatment showed significantly reduced ovarian and body weights of PCOS rats. Farzana *et al.* [35] showed a dignified diminution in ovarian weight and follicles number after flaxseed administration in PCOS patients without any change in body weight while Pareek & Boolchandani [36] revealed a decrease in the mean of body mass index with no change in body weight. Cystic follicles number and thickness of the theca layer significantly decreased after flaxseed administration in EV-induced rats [37]. However, both plants containing compounds that benefit these two parameters by many means such as lignans, polyunsaturated fatty acids, alkaloids, steroids, flavonoids, and saponins [32-40].

EV induction caused a rise in blood glucose, insulin, and IR, which was consistent with Sasikala *et al.* [41], EV supplementation caused high androgen and IR leading to hyperinsulinemia. Several studies reported abnormal glucose metabolism associated with PCOS [42–44] and that might mainly be due to IR present in this syndrome. Fenugreek aqueous extract exhibited a considerable decline in serum glucose and insulin and improved IR, proving its hypoglycemic effect. This effect may be mediated by three ways, lowering glucose, decreasing cells sensitivity to insulin or by exhibiting insulin-like effects. Fibers in fenugreek play the main role in its hypoglycemic effect, active components and phytochemicals such galactomannan, [45] mycelin, flavonoids, saponins, glucosides, and pectin have shown antidiabetic effects that differ in the way of reducing glucose level [46]. Some components in fenugreek are also found in flaxseed including fiber and its glucose-lowering effect. There are different results about the effect of flaxseed on the glucose level. For example, Javidi *et al.* [47] observed no change in glucose and IR, when administered flaxseed powder daily to prediabetes patients, while Pan *et al.* [48] reported small but significant improvement in glycemic control in T2DM patients after lignan supplementation with no change in glucose and lipid profile or IR. In accordance with our study, Thakur *et al.* [49] observed a decrease in blood glucose after flaxseed gum supplementation in type II diabetic persons. Recently, Kapuriya *et al.* [24] announced that the flaxseed aqueous extract has antidiabetic activity in an experimental rat model. The current study showed that the combination of fenugreek and flaxseed possess the highest reduction effect on glucose among treatment groups.

In the current results, EV produced a significant rise in serum TC, TG, and LDL-c along with a significant drop in HDL-c. Knauff *et al.* [50] found that reduced insulin sensitivity involved in high TG concentration is associated with an increase in free androgens in PCOS. Since IR neglects insulin function in PCOS, hyperinsulinemia occurs; high lipid profile is expected in PCOS. In agreement with our results, Macut *et al.* [51], Lath [52], and Iuhās *et al.* [53] studies revealed that serum insulin, IR, TC, TG, and LDL-c levels were greater in PCOS compared to normal except HDL-c that was lower. Treatment of EV-induced rats with fenugreek and flaxseed aqueous extracts was notably diminished serum TC, TG, and LDL-c and raised HDL-c compared to untreated rats. Numerous studies showed that the same active components of fenugreek, which are involved in glucose reduction also participate in its lowering effect on lipids. Kassaian *et al.* [54] and Gupta & Verma [55] findings showed that administration of fenugreek to type II diabetic patients lowered TC, TG, VLDL-C, and LDL and lifted up HDL level, which might be due to the saponin and galactomannan of fenugreek that reduces its absorption from the intestine. lipid-lowering effect of fenugreek is owing to its suppression of fast assemblage and upregulation of low-density lipoprotein receptor [56]. Treating PCOS-induced rats with flaxseed showed a result similar to fenugreek and lipid profile was recovered. Thakur *et al.* [49] found a decrease in TC and LDL-c while Devarshi *et al.* [57] found a decrease in TG and VLDL-c with an increase in HDL in diabetic subjects after flaxseed supplementation. Torkan *et al.* [58] reported that flaxseed

administration to hyperlipidemic individuals reduced TC, LDL, and TG along with body weight and mass index. Kristensen *et al.* [59] showed that flaxseed dietary fibers decreased TC and LDL-c and increased fat excretion while Newairy & Abdou [60] revealed that reduced lipids were attributed to protective effects of lignan isolated from flaxseed. Abd El Ghany *et al.* [61] concluded that the lipid-lowering effect of flaxseed might be as a result of their rich content of polyphenols and flavonoids.

No change in estradiol but significant change in other measured reproductive hormones parameters was observed in PCOS rats; testosterone, FSH, and LH increased while progesterone decreased. In semi-agreement with our results, Pournaderi *et al.*[62] found a significantly high level of FSH, LH, and testosterone as well as E2 and progesterone in PCOS-induced subjects in contrast to control. Kakadia *et al.*[63] and Li *et al.*[64] as well found increased LH, FSH, and T while reduced E2 and progesterone in PCOS Letrozole-induced group. A study on Saudi PCOS women showed increased level of T and LH while FSH, progesterone and sex hormone-binding globulin (SHBG) levels decreased[65]. High insulin might be involved in high androgen. Botwood *et al.* [66] revealed that SHBG synthesis and secretion are inhibited by insulin. In this trend, Braga *et al.* [67] and Malini & Roy George [68] found that IR, insulin, T, and LH were high in PCOS women while SHBG was lower. Novak *et al.* [69] proposed that PCOS anovulatory condition is due to high LH level caused by the abnormal passive feedback of E2 or progesterone on LH excretion. Because of various complications of PCOS and its unclear origin, it is difficult to point the exact mechanism involved in the syndrome. Fenugreek treatment showed a significant restoration in all of the measured sex hormones in EV-induced rats. Modaresi *et al.* [70] showed that fenugreek extract decreased FSH and LH and increased E2 along with folliculogenesis reduction in ovaries and concluded that fenugreek has infertility effect on healthy female mice. Swaroop *et al.* [30] showed that fenugreek reduced LH/FSH ratio and the number of ovarian cysts and also improved menstrual cycle in women with PCOS and some got pregnant. It is important to mention that fenugreek has a lipid-lowering effect and since E2 and androgen are mainly made of dehydroepiandrosterone (DHEA), which is made from cholesterol [71], it might help in androgen reduction. EV-induced rats treated with flaxseed revealed results similar to fenugreek so that T, FSH, and LH increased and progesterone decreased in them. As discussed, flaxseed enriched with lignans showed androgen reduction effect in PCOS women [72]. Jelodar *et al.* [37] observed an increase in progesterone level with a decrease in T after flaxseed extract supplementation to PCOS-induced rats, with no change in E2 and DHEA. Rietjens *et al.* [73] demonstrated that phytoestrogens have a structure similar to E2 enabling them to provoke anti-estrogenic effect by binding to estrogen receptors. Barnes *et al.* [74] as well mentioned that estrogenic or anti-estrogenic effect of phytoestrogen depends on the endogenous estrogen concentrations. Low fat intake might reduce androgen level, which might be due to flaxseed fibers and their decreasing effect on cholesterol. We think that reduction in androgen might be related first, to lipid reduction effect of flaxseed fibers, and second, to flaxseed lignans metabolites that may release anti-androgenic effect attributed to their negative feedback and reducing testosterone synthesis.

The liver has a vital executive role in various metabolic procedures, including lipid synthesis, transmission, and storage, as well as glucose and insulin metabolism that all related to PCOS [75]. Our result showed a significant increase in ALT, AST in PCOS group indicating the appearance of disrupted hepatic function in PCOS. Karoli *et al.* [76] and Gambarin-Gelwan *et al.* [77] detected similar results of increased liver enzymes in PCOS women, which is related to IR, obesity, dyslipidemia, and aging. Cree-Green *et al.* [78], Minato *et al.* [3], and Schwimmer *et al.* [79] findings showed an increase in AST and observed high IR, BMI, and dyslipidemia in PCOS subjects. In addition to high AST level, ALT and alkaline phosphate were high in EV-induced rats. Supplementation of fenugreek and/or flaxseed to EV-induced group manifested a notably reduced in the measured liver enzymes. El-Wakf *et al.* [80] observed a significant reduction in serum ALT and AST in nitrate-exposed rats after fenugreek administration. Moreover, Kaviarasan & Anuradha [81] used the polyphenolic extract of fenugreek seed to decrease liver damage and the increased liver enzymes in rats with chronic ethanol administration, which were retrieved after extract administration. Kandhare *et al.* [82] confirmed the existence of anti-inflammatory, antioxidants, and anti-fibrotic activities in fenugreek extract given to induced-liver fibrosis rats with a decrease in AST, ALT, and GGT. This may be caused by phytochemicals in fenugreek. Yari *et al.* [83] showed the hepatoprotective effect of flaxseed against non-alcoholic fatty liver disease (NAFLD) in patients who administered it, causing reduced liver enzyme, IR, body weight, hepatic fibrosis, and fatty liver disease. Al-Bishri [84] said that flaxseed polyunsaturated fatty acids (PUFAs) comprising omega-3 and omega-6 fatty acids are responsible for the reduction of liver markers in hypertensive rats.

The ovarian of EV-induced rats showed oxidative stress (OS) sign, in which oxidants were higher than antioxidants. Birben *et al.* [85] implied that many disorders showed OS markers and reactive oxygen species

(ROS) produced from cellular metabolic activities and environmental factors. In PCOS, IR provokes OS production due to hyperglycemia and hyperlipidemia leading to ROS production [85,86]. Ghowsi *et al.* [86] evidenced that androgen is related to excess LPO and other OS markers in PCOS-induced rats. Papalou *et al.* [87] considered that PCOS is mainly an oxidative state where antioxidants unable to scavenge the excess free radicals. Some studies showed a significant decrease in SOD [88] and GSH while an increase in MDA and NO [89, 90] in PCOS subjects. Treating PCOS-induced rats with fenugreek and flaxseed have attenuated the imbalanced antioxidants to oxidants. Many studies showed the antioxidant effect of fenugreek against OS in PCOS and other disorders [91, 92]. Herrera *et al.* [93] and Baba *et al.* [94] detected a high amount of flavonoids and saponins in fenugreek extract. Subhashini & *et al.* [95], Dixit *et al.* [96], Mohammadzadeh *et al.* [97], Pandey & Awasthi [98] and Abeed Al Mashkor [99] showed that polyphenols and flavonoids are responsible for fenugreek antioxidant effect and scavenging free radicals in many disorders. Flaxseed also showed a protective effect against OS in PCOS and many other diseases [100–103]. Concerning flaxseed, various studies showed that lignans mainly secoisolariciresinol diglucoside (SDG) in flaxseed responsible in its antioxidant activity [60,104,105] as well as PUFAs such as omega-3 fatty acid [103] and rich phenolic compounds[104].

## CONCLUSION:

The current study focused on the effect of fenugreek on insulin and IR and its hypoglycemic effect in PCOS-induced rats, as well as anti-androgenic effect of flaxseed. Both aqueous extracts of the two plants showed beneficial effect against serum glucose, insulin, IR, lipid profile, reproductive hormones, liver function, and OS. The combination of both fenugreek and flaxseed extracts showed better amelioration on all of these parameters and work synergistically to achieve the expected results. Therefore, current and future investigations will contribute to bridging the traditional knowledge of using natural products combination to reduce the symptoms of PCOS.

## REFERENCES

1. Subhi Ramadhan, R., Molecular analysis of FSH receptor gene in Iraqi women with PCOS syndrome, Middle East Fertil. Soc. J., 2018, 23(4), 404–408
2. Abou-El-Naga, M., Habbak, Z., El Bakary, E.R., Potential effects of Mentha piperita ( peppermint ) on Letrozole- induced polycystic ovarian syndrome in female albino rat, Int. J. Adv. Res., 2015, 3(10), 211–226
3. Minato, S., Sakane, N., Kotani, K., Nirengi, S., et al., Prevalence and Risk Factors of Elevated Liver Enzymes in Japanese Women With Polycystic Ovary Syndrome., J. Clin. Med. Res., 2018, 10(12), 904–910
4. Heibashy, M.I.A., Mazen, G.M.A., Shahin, M.I., Metabolic Changes and Hormonal Disturbances in Polycystic Ovarian Syndrome Rats and the Amelioration Effects of Metformin and/or Cinnamon Extraction, Journa Am. Sci., 2013, 9(12), 54–62
5. Yilmaz, B., Vellanki, P., Ata, B., Yildiz, B.O., Metabolic syndrome, hypertension, and hyperlipidemia in mothers, fathers, sisters, and brothers of women with polycystic ovary syndrome: a systematic review and meta-analysis, Fertil. Steril., 2018, 109(2), 356–364
6. Tripathy, P., Sahu, A., Sahu, M., Nagy, A., Metabolic risk assessment of Indian women with polycystic ovarian syndrome in relation to four Rotterdam criteria based phenotypes, Eur. J. Obstet. Gynecol. Reprod. Biol., 2018, 224, 60–65
7. Ning, N., Balen, A., Brezina, P.R., Leong, M., et al., How to recognize PCOS: Results of a web-based survey at IVF-worldwide.com, Reprod. Biomed. Online, 2013, 26(5), 500–505
8. Akgül, S., Düzçeker, Y., Kanbur, N., Derman, O., Do Different Diagnostic Criteria Impact Polycystic Ovary Syndrome Diagnosis for Adolescents?, J. Pediatr. Adolesc. Gynecol., 2018, 31(3), 258–262
9. Allahbadia, G.N., Merchant, R., Polycystic ovary syndrome and impact on health, Middle East Fertil. Soc. J., 2011, 16(1), 19–37
10. Paixão, L., Ramos, R.B., Lavarda, A., Morsh, D.M., Spritzer, P.M., Animal models of hyperandrogenism and ovarian morphology changes as features of polycystic ovary syndrome: a systematic review., Reprod. Biol. Endocrinol., 2017, 15(1), 12
11. Mansour, A., Hosseini, S., Larijani, B., Mohajeri-Tehrani, M.R., Nutrients as novel therapeutic approaches for metabolic disturbances in polycystic ovary syndrome., EXCLI J., 2016, 15, 551–564



12. Patel, R., Shah, G., Insulin Sensitizers Modulate GnRH Receptor Expression in PCOS Rats, Arch. Med. Res., 2018, 49(3), 154–163
13. Sirmans, S.M., Pate, K.A., Epidemiology, diagnosis, and management of polycystic ovary syndrome, Clin. Epidemiol., 2013, 6(1), 1–13
14. Yıldırım, D.İ., Marakoğlu, K., Complementary and alternative medicine use amongst Turkish type 2 diabetic patients: A cross-sectional study, Complement. Ther. Med., 2018, 41, 41–46
15. Saeed S, Islahudin F, Makmor-Bakry M, Redzuan AM. The practice of complementary and alternative medicine among chronic Kidney disease patients. Journal of Advanced Pharmacy Education & Research| Jul-Sep. 2018 Jan 1;8(3):31.
16. Scarton, L.A., Wang, L., Kilicoglu, H., Jahries, M., Del Fiol, G., Expanding vocabularies for complementary and alternative medicine therapies, Int. J. Med. Inform., 2018, 121, 64–74
17. Poole, C., Bushey, B., Foster, C., Campbell, B., et al., The effects of a commercially available botanical supplement on strength, body composition, power output, and hormonal profiles in resistance-trained males., J. Int. Soc. Sports Nutr., 2010, 7(1), 34
18. Al-Dabbagh, B., Elhaty, I.A., Al Hrou, A., Al Sakkaf, R., et al., Antioxidant and anticancer activities of Trigonella foenum-graecum, Cassia acutifolia and Rhazya stricta, BMC Complement. Altern. Med., 2018, 18(1), 240
19. Baliga, M.S., Palatty, P.L., Adnan, M., Naik, T.S., et al., Anti-Diabetic Effects of Leaves of Trigonella foenum-graecum L. (Fenugreek): Leads from Preclinical Studies, J. Food Chem. Nanotechnol., 2017, 03(02), 67–71
20. Abdelatif, A.M., Ibrahim, M.Y., Mahmoud, A.S., Antidiabetic Effects of Fenugreek (Trigonella foenum-graecum) Seeds in the Domestic Rabbit (*Oryctolagus cuniculus*), Res. J. Med. Plant, 2012, 6(6), 449–455
21. Pundarikakshudu, K., Shah, D., Panchal, A., Bhavsar, G., Anti-inflammatory activity of fenugreek (Trigonella foenum-graecum Linn) seed petroleum ether extract, Indian J. Pharmacol., 2016, 48(4), 441
22. Braun, L., Cohen, M., Herbs & natural supplements: An Evidence based Guide, Elsevier Australia, 2010,
23. Bernacchia, R., Preti, R., Vinci, G., Chemical Composition and Health Benefits of Flaxseed, Austin J. Nutr. Food Sci., 2014, 2(8), 1–9
24. Kapuriya, P.B., Bhavsar, S.K., Thaker, A.M., Sadariya, K.A., Antidiabetic activity of aqueous extracts of *Linum usitatissimum* in streptozotocin induced diabetic rats, Pharma Innov. J., 2018, 7(7), 149–154
25. Rashed, M.M., Shallan, M., Mohamed, D.A., Fouda, K., Hanna, L.M., Biological Evaluation of Anti-androgenic Effect of Some Plant Foods, J. Food Nutr. Res., 2014, 2(9), 645–651
26. Hasan, M., Rahman, M., Effect of Fenugreek on Type2 diabetic patients, Int. J. Sci. Res. Publ., 2016, 6(1), 251–2250
27. Ghafurniyan, H., Azarnia, M., Nabiuni, M., Karimzadeh, L., The effect of green tea extract on reproductive improvement in estradiol valerate-induced polycystic ovary polycystic ovarian syndrome in rat, Iran. J. Pharm. Res., 2015, 14(4), 1215–1223
28. Karimzadeh, L., Nabiuni, M., Kouchesfehiani, H., Adham, H., et al., Effect of bee venom on IL-6, COX-2 and VEGF levels in polycystic ovarian syndrome induced in Wistar rats by estradiol valerate, J. Venom. Anim. Toxins Incl. Trop. Dis., 2013, 19(1), 32
29. Jashni, H.K., Jahromi, H.K., Bagheri, Z., The Effect of Palm Pollen Extract on Polycystic Ovary Syndrome (POS) in Rats, International J. Med. Res. Heal. Sci., 2016, 5(S), 317–321
30. Swaroop, A., Jaipuria, A.S., Gupta, S.K., Bagchi, M., et al., Efficacy of a Novel Fenugreek Seed Extract (Trigonella foenum-graecum, Furocyst) in Polycystic Ovary Syndrome (PCOS), Int. J. Med. Sci., 2015, 12(10), 825–831
31. Muraki, E., Chiba, H., Taketani, K., Hoshino, S., et al., Fenugreek with reduced bitterness prevents diet-induced metabolic disorders in rats, Lipids Health Dis., 2012, 11, 58
32. Moradi, N., Didarshetaban, M.B., Reza, H., Pour, S., Fenugreek (Trigonella foenum-graecum L.) As a Valuable Medicinal Plant, Int. J. Adv. Biol. Biomed. Res., 2013, 1(8), 922–931
33. Maha A. Althaiban. Antiulcer potential of olive leaves extract in gastric ulcer induced by indomethacin in male rats: antioxidant and anti-inflammatory effects. Pharmacophore. 2018.9(6):57-64.
34. Tabet A, Boukhari A, Nouidjem Y. Phenolic content, HPLC analysis and Antioxidant activity extract from Tamarix Articulata. Journal of Advanced Pharmacy Education & Research| Oct-Dec. 2018;8(4).
35. Farzana, F., Sulaiman, A.F., Ruckmani, A., Vijayalakshmi, K., et al., Effects of Flax Seeds Supplementation in PolyCystic Ovarian Syndrome 2015, 31(23), 113–119

36. Pareek, M., Boolchandani, R., Effect of Flaxseed Supplementation on Hormone Profile in Women with PCOS, *International J. Food, Agric. Vet. Sci.*, 2018, 8(1), 19–25
37. Jelodar, G., Masoomi, S., Rahmanifar, F., Hydroalcoholic extract of flaxseed improves polycystic ovary syndrome in a rat model., *Iran. J. Basic Med. Sci.*, 2018, 21(6), 645–650
38. Rahmani, E., Samimi, M., Ebrahimi, F.A., Foroozanfard, F., et al., The effects of omega-3 fatty acids and vitamin E co-supplementation on gene expression of lipoprotein(a) and oxidized low-density lipoprotein, lipid profiles and biomarkers of oxidative stress in patients with polycystic ovary syndrome, *Mol. Cell. Endocrinol.*, 2017, 439, 247–255
39. Vargas, M.L., Almario, R.U., Buchan, W., Kim, K., Karakas, S.E., Metabolic and endocrine effects of long-chain versus essential omega-3 polyunsaturated fatty acids in polycystic ovary syndrome, *Metabolism*, 2011, 60(12), 1711–1718
40. Moree, S.S., Secoisolariciresinol Diglucoside: A potent multifarious bioactive phytoestrogen of flaxseed, *Res. Rev. Biomed. Biotechnol.*, 2011, 2(323), 1–24
41. Sasikala, M., Parthiban, P., Vijayakumar, S., A Comprehensive Review on Polycystic Ovary Syndrome and Its Therapeutic Management For Ovulation Induction In Infertile Women-Part-I, *Int. J. Adv. Case Reports*, 2014, 1(2), 64–77
42. Teede, H., Deeks, A., Moran, L., Polycystic ovary syndrome: a complex condition with psychological, reproductive and metabolic manifestations that impacts on health across the lifespan, *BMC Med.*, 2010, 8(1), 41
43. Dumitrescu, R., Mehedintu, C., Briceag, I., Purcarea, V.L., Hudita, D., The polycystic ovary syndrome: An update on metabolic and hormonal mechanisms, *J. Med. Life*, 2015, 8(2), 142–145
44. Li, S., Chu, Q., Ma, J., Sun, Y., et al., Discovery of novel lipid profiles in pcos: Do insulin and androgen oppositely regulate bioactive lipid production?, *J. Clin. Endocrinol. Metab.*, 2017, 102(3), 810–821
45. Srichamroen, A., Thomson, A.B.R., Field, C.J., Basu, T.K., In vitro intestinal glucose uptake is inhibited by galactomannan from Canadian fenugreek seed (*Trigonella foenum graecum* L) in genetically lean and obese rats, *Nutr. Res.*, 2009, 29(1), 49–54
46. Gothai, S., Ganesan, P., Park, S.Y., Fakurazi, S., et al., Natural phyto-bioactive compounds for the treatment of type 2 diabetes: Inflammation as a target, *Nutrients*, 2016, 8(8), 461
47. Javidi, A., Mozaffari-Khosravi, H., Nadjarzadeh, A., Dehghani, A., Eftekhari, M.H., The effect of flaxseed powder on insulin resistance indices and blood pressure in prediabetic individuals: A randomized controlled clinical trial, *J. Res. Med. Sci.*, 2016, 21(5), 70
48. Pan, A., Sun, J., Chen, Y., Ye, X., et al., Effects of a Flaxseed-Derived Lignan Supplement in Type 2 Diabetic Patients: A Randomized, Double-Blind, Cross-Over Trial, *PLoS One*, 2007, 2(11), e1148
49. Thakur, G., Mitra, A., Pal, K., Rousseau, D., Effect of flaxseed gum on reduction of blood glucose and cholesterol in type 2 diabetic patients, *Int. J. Food Sci. Nutr.*, 2009, 60(sup6), 126–136
50. Knauff, E.A.H., Westerveld, H.E., Goverde, A.J., Eijkemans, M.J., et al., Lipid profile of women with premature ovarian failure, *Menopause*, 2008, 15(5), 919–923
51. Macut, D., Panidis, D., Glišić, B., Spanos, N., et al., Lipid and lipoprotein profile in women with polycystic ovary syndrome This article is one of a selection of papers published in the special issue Bridging the Gap: Where Progress in Cardiovascular and Neurophysiologic Research Meet., *Can. J. Physiol. Pharmacol.*, 2008, 86(4), 199–204
52. Lath, R., Insulin resistance and lipid profile in polycystic ovary syndrome, *Asian J. Biomed. Pharm. Sci.*, 2015, 5(47), 30–35
53. Iuhas, C., Costin, N., Miha, D., Lipid Parameters in Patients with Polycystic Ovary Syndrome, *Appl. Med. Informatics*, 2012, 31(4), 27–32
54. Kassaian, N., Azadbakht, L., Forghani, B., Amini, M., Effect of Fenugreek Seeds on Blood Glucose and Lipid Profiles in Type 2 Diabetic Patients, *Int. J. Vitam. Nutr. Res.*, 2009, 79(1), 34–39
55. Gupta, R., Verma, S., Study of Effects of Consumption of Powdered Fenugreek Seeds on Serum Lipid Profile in Patients with Type 2 Diabetes Mellitus, *J. Pak. Med. Assoc.*, 2015, 6(1), 57–59
56. Vijayakumar, M., Pandey, V., Mishra, G.C., Bhat, M.K., Hypolipidemic effect of fenugreek seeds is mediated through inhibition of fat accumulation and upregulation of LDL receptor., *Obesity (Silver Spring)*, 2010, 18(4), 667–674

57. Devarshi, P.P., Jangale, N.M., Ghule, A.E., Bodhankar, S.L., Harsulkar, A.M., Beneficial effects of flaxseed oil and fish oil diet are through modulation of different hepatic genes involved in lipid metabolism in streptozotocin-nicotinamide induced diabetic rats., *Genes Nutr.*, 2013, 8(3), 329–42
58. Torkan, M., Entezari, M.H., Siavash, M., Effect of flaxseed on blood lipid level in hyperlipidemic patients., *Rev. Recent Clin. Trials*, 2015, 10(1), 61–7
59. Kristensen, M., Jensen, M.G., Aarestrup, J., Petersen, K.E., et al., Flaxseed dietary fibers lower cholesterol and increase fecal fat excretion, but magnitude of effect depend on food type., *Nutr. Metab. (Lond.)*, 2012, 9, 1–8
60. Newairy, A.S.A., Abdou, H.M., Protective role of flax lignans against lead acetate induced oxidative damage and hyperlipidemia in rats, *Food Chem. Toxicol.*, 2009, 47(4), 813–818
61. Abd El Ghany, M., EL Mehiry, H.F., Shelbaya, L.A., Reziq, A.A., Potential Effect of Dietary Flaxseed (*Linum Usitatissimum* L.) Powder and Extract on aged Menopausal Female Rats, *IOSR J. Environ. Sci. Toxicol. Food Technol.*, 2017, 11(9), 2319–2399
62. Pournaderi, P.S., Yaghmaei, P., Khodaei, H., Noormohammadi, Z., Hejazi, S.H., The effects of 6-Gingerol on reproductive improvement, liver functioning and Cyclooxygenase-2 gene expression in estradiol valerate – Induced polycystic ovary syndrome in Wistar rats, *Biochem. Biophys. Res. Commun.*, 2017, 484(2), 461–466
63. Kakadia, N., Patel, P., Deshpande, S., Shah, G., Effect of *Vitex negundo* L. seeds in letrozole induced polycystic ovarian syndrome, *J. Tradit. Complement. Med.*, 2018, 1–10
64. Li, C., Chen, L., Zhao, Y., Chen, S., et al., Altered expression of miRNAs in the uterus from a letrozole-induced rat PCOS model, *Gene*, 2017, 598, 20–26
65. Fakhoury, H., Tamim, H., Ferwana, M., Siddiqui, I., et al., Age and BMI Adjusted Comparison of Reproductive Hormones in PCOS, *J. Fam. Med. Prim. Care*, 2012, 1(2), 132
66. Botwood, N., Hamilton-Fairley, D., Kiddy, D., Robinson, S., Franks, S., Sex hormone-binding globulin and female reproductive function., *J. Steroid Biochem. Mol. Biol.*, 1995, 53(1–6), 529–31
67. Braga, L.D. da C., Godoy-Matos, A.F., de Oliveira Siciliano, P., Corrêa do Amaral, J.O., Carvalho, D.P., Is DPP4 activity increased in PCOS?, *Diabetes Metab. Syndr. Clin. Res. Rev.*, 2018, 12(5), 673–675
68. Malini, N.A., Roy George, K., Evaluation of different ranges of LH:FSH ratios in polycystic ovarian syndrome (PCOS) – Clinical based case control study, *Gen. Comp. Endocrinol.*, 2018, 260, 51–57
69. Novak, I., Janeiro, P., Seruga, M., Oliveira-Brett, A.M., Ultrasound extracted flavonoids from four varieties of Portuguese red grape skins determined by reverse-phase high-performance liquid chromatography with electrochemical detection, *Anal. Chim. Acta*, 2008, 630(2), 107–115
70. Modaresi, M., Mahdian, B., Jalalizand, A., in: *Int. Conf. Appl. Life Sci.*, 2012, pp. 437–443
71. Miller, W.L., Disorders of androgen synthesis - From cholesterol to dehydroepiandrosterone, *Med. Princ. Pract.*, 2005, 14(1), 58–68
72. Nowak, D.A., Snyder, D.C., Brown, A.J., Demark-Wahnefried, W., The effect of flaxseed supplementation on hormonal levels associated with polycystic ovarian syndrome: A case study, *Curr. Top. Nutraceutical Res.*, 2007, 5(4), 177–181
73. Rietjens, I.M.C.M., Lousse, J., Beekmann, K., The potential health effects of dietary phytoestrogens, *Br. J. Pharmacol.*, 2017, 174(11), 1263–1280
74. Barnes, S., Peterson, T.G., Coward, L., Rationale for the use of genistein-containing soy matrices in chemoprevention trials for breast and prostate cancer, *J. Cell. Biochem.*, 1995, 59(S22), 181–187
75. Paschos, P., Paletas, K., Non alcoholic fatty liver disease and metabolic syndrome, *Hippokratia*, 2009, 13(1), 9–19
76. Karoli, R., Fatima, J., Chandra, A., Gupta, U., et al., Prevalence of hepatic steatosis in women with polycystic ovary syndrome., *J. Hum. Reprod. Sci.*, 2013, 6(1), 9–14
77. Gambarin–Gelwan, M., Kinkhabwala, S. V., Schiano, T.D., Bodian, C., et al., Prevalence of Nonalcoholic Fatty Liver Disease in Women With Polycystic Ovary Syndrome, *Clin. Gastroenterol. Hepatol.*, 2007, 5(4), 496–501
78. Cree-Green, M., Bergman, B.C., Coe, G. V, Newnes, L., et al., Hepatic Steatosis is Common in Adolescents with Obesity and PCOS and Relates to De Novo Lipogenesis but not Insulin Resistance, *Obesity*, 2016, 24(11), 2399–2406
79. Schwimmer, J.B., Khorram, O., Chiu, V., Schwimmer, W.B., Abnormal aminotransferase activity in women with polycystic ovary syndrome, *Fertil. Steril.*, 2005, 83(2), 494–497

80. El-Wakf, A.M., Hassan, H.A., Mahmoud, A.Z., Habza, M.N., Fenugreek potent activity against nitrate-induced diabetes in young and adult male rats, *Cytotechnology*, 2015, 67(3), 437–447
81. Kaviarasan, S., Anuradha, C. V, Fenugreek (*Trigonella foenum graecum*) seed polyphenols protect liver from alcohol toxicity: a role on hepatic detoxification system and apoptosis., *Pharmazie*, 2007, 62(4), 299–304
82. Kandhare, A.D., Bodhankar, S.L., Mohan, V., Thakurdesai, P.A., Glycosides Based Standardized Fenugreek Seed Extract Ameliorates Bleomycin-induced Liver Fibrosis in Rats Via Modulation of Endogenous Enzymes., *J. Pharm. Bioallied Sci.*, 2017, 9(3), 185–194
83. Yari, Z., Rahimlou, M., Eslamparast, T., Ebrahimi-Daryani, N., et al., Flaxseed supplementation in non-alcoholic fatty liver disease: a pilot randomized, open labeled, controlled study, *Int. J. Food Sci. Nutr.*, 2016, 67(4), 461–469
84. Al-Bishri, W.M., Favorable effects of flaxseed supplemented diet on liver and kidney functions in hypertensive Wistar rats., *J. Oleo Sci.*, 2013, 62(9), 709–15
85. Birben, E., Sahiner, U.M., Sackesen, C., Erzurum, S., Kalayci, O., Oxidative stress and antioxidant defense., *World Allergy Organ. J.*, 2012, 5(1), 9–19
86. Ghowsi, M., Khazali, H., Sisakhtnezhad, S., The effect of resveratrol on oxidative stress in the liver and serum of a rat model of polycystic ovary syndrome: An experimental study., *Int. J. Reprod. Biomed.*, 2018, 16(3), 149–158
87. Papalou, O., Victor, V.M., Diamanti-Kandarakis, E., Oxidative Stress in Polycystic Ovary Syndrome., *Curr. Pharm. Des.*, 2016, 22(18), 2709–2722
88. Seleem, A.K., El Refaeey, A.A., Shaalan, D., Sherbiny, Y., Badawy, A., Superoxide dismutase in polycystic ovary syndrome patients undergoing intracytoplasmic sperm injection., *J. Assist. Reprod. Genet.*, 2014, 31(4), 499–504
89. Karabulut, A., Cakmak, M., Kiran, R., Sahin, I., Oxidative Stress Status, Metabolic Profile and Cardiovascular Risk Factors in Patients with Polycystic Ovary Syndrome, *Med. Sci.*, 2012, 1(1), 27–34
90. Murri, M., Luque-ramírez, M., Insenser, M., Ojeda-ojeda, M., Escobar-morreale, H.F., Circulating markers of oxidative stress and polycystic ovary syndrome (pcos): A systematic review and meta-analysis, *Hum. Reprod. Update*, 2013, 19(3), 268–288
91. Herrera, T., Navarro del Hierro, J., Fornari, T., Reglero, G., Martin, D., Inhibitory effect of quinoa and fenugreek extracts on pancreatic lipase and  $\alpha$ -amylase under in vitro traditional conditions or intestinal simulated conditions, *Food Chem.*, 2019, 270, 509–517
92. Baba, W.N., Tabasum, Q., Muzzaffar, S., Masoodi, F.A., et al., Some nutraceutical properties of fenugreek seeds and shoots (*Trigonella foenum-graecum* L.) from the high Himalayan region, *Food Biosci.*, 2018, 23, 31–37
93. Subhashini, N., Thangathirupathi, A., Lavanya, N., Antioxidant Activity of *Trigonella Foenum Graecum* Using Various in Vitro and Ex Vivo Models, *Int J Pharm Pharm Sci*, 2011, 3(2), 96–102
94. Dixit, P., Ghaskadbi, S., Mohan, H., Devasagayam, T.P.A., Antioxidant properties of germinated fenugreek seeds, *Phyther. Res.*, 2005, 19(11), 977–983
95. Mohammadzadeh, A., Gol, A., Oloumi, H., The Effects of Fenugreek Seed Powder on Oxidant and Antioxidant Factors in Male Rats with Acetaminophen-induced Liver Toxicity, *J Babol Univ Med Sci*, 2015, 17(9), 44–51
96. Pandey, H., Awasthi, P., Effect of processing techniques on nutritional composition and antioxidant activity of fenugreek (*Trigonella foenum-graecum*) seed flour., *J. Food Sci. Technol.*, 2015, 52(2), 1054–60
97. Abeed Al Mashkor, I.M., Phenolic Content and Antioxidant Activity of Fenugreek Seeds Extract, *Int. J. Pharmacogn. Phytochem. Res.*, 2014, 6(4), 841–844
98. Badawy, E., Rasheed, W., Elias, T., Hussein, J., et al., Flaxseed oil reduces oxidative stress and enhances brain monoamines release in streptozotocin-induced diabetic rats, *Hum. Exp. Toxicol.*, 2015, 34(11), 1133–1138
99. Abdel-Moniem, A.E., Dkhil, M.A., Al-Quraishy, S., Protective role of flaxseed oil against lead acetate induced oxidative stress in testes of adult rats, *African J. Biotechnol.*, 2010, 9(42), 7216–7223
100. Rizwan, S., Naqshbandi, A., Farooqui, Z., Khan, A.A., Khan, F., Protective effect of dietary flaxseed oil on arsenic-induced nephrotoxicity and oxidative damage in rat kidney, *Food Chem. Toxicol.*, 2014, 68, 99–107

101. Wahba, H.M.A., Ibrahim, T.A.A., Protective effect of flaxseed oil and vitamin E on potassium Bromate-induced oxidative stress in male rats, *Int J Curr Microbiol App Sci*, 2013, 2(9), 299–309
102. Prasad, K., Regression of hypercholesterolemic atherosclerosis in rabbits by secoisolariciresinol diglucoside isolated from flaxseed, *Atherosclerosis*, 2008, 197(1), 34–42
103. Pilar, B., Güllich, A., Oliveira, P., Ströher, D., et al., Protective Role of Flaxseed Oil and Flaxseed Lignan Secoisolariciresinol Diglucoside Against Oxidative Stress in Rats with Metabolic Syndrome, *J. Food Sci.*, 2017, 82(12), 3029–3036
104. Austria, J.A., Richard, M.N., Chahine, M.N., Edel, A.L., et al., Bioavailability of alpha-linolenic acid in subjects after ingestion of three different forms of flaxseed., *J. Am. Coll. Nutr.*, 2008, 27(2), 214–21
105. Kaur, R., Kaur, M., Microstructural, physicochemical, antioxidant, textural and quality characteristics of wheat muffins as influenced by partial replacement with ground flaxseed, *LWT*, 2018, 91, 278–285.