



Research Article

ISSN : 2277-3657
CODEN(USA) : IJPRPM

Reduction of High Glucose-Induced Nitric Oxide Synthase Expression in Human Vascular Endothelial Cells by Ascorbic Acid and α -Tocopherol

Mohammad Ali Ghaffari¹, Mohammad Aberumand², Hoshang Roshanmehr²
and Seyed Ahmad Hosseini^{*3}

¹Cellular and Molecular Research Center, Ahvaz Jundishapur, University of Medical Sciences, Ahvaz, Iran

²Department of Biochemistry, School of Medical, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran,

³Nutrition and Metabolic Diseases Research Center, Ahvaz Jundishapur, University of Medical Sciences, Ahvaz, Iran

*Email: seyedahmadhosseini@yahoo.com

ABSTRACT

Hyperglycemia is generally regarded as one of the major causes of vascular complications in diabetic patients. It has been shown that high-glucose concentration increases nitric oxide synthase (NOS) expression and NO generation in cultured human aortic endothelial cells. The epidemiological studies have demonstrated that there is an inverse relation between vitamin use and cardiovascular disease in diabetes patients. This study tested whether α -tocopherol and ascorbic acid can alter high glucose-induced expression of NOS in vascular endothelial cells. Human umbilical vascular endothelial cells (HUVECs) were treated with α -tocopherol and ascorbic acid, and stimulated by high glucose. Expression of endothelial nitric oxide synthase (eNOS) and inducible nitric oxide synthase (iNOS) was measured by real-time PCR (RT-PCR). Ascorbic acid (100 μ M) and α -tocopherol (200 μ M) significantly inhibited glucose (25 mM)-induced expression of eNOS mRNA and iNOS mRNA. Ascorbic acid and α -tocopherol are also able to reduce concentration of NOS in HUVECs. Additionally, there is a significant positive relation between expression of eNOS and iNOS mRNA with concentration of NOS. These results suggest that ascorbic acid and α -tocopherol can inhibit high glucose-induced production of NOS proteins by downregulation of its gene expression. Therefore, important role of ascorbic acid and α -tocopherol in the treatment of vascular dysfunction associated with diabetes disease may be via this molecular mechanism. Of course, further studies on regulation of these vitamins to signaling pathways are necessary.

Key words: Glucose, HUVECs, nitric oxide synthase, α -tocopherol, ascorbic acid

INTRODUCTION

Diabetes mellitus is a frequent and increasing public health problem. Diabetes has a prevalence of 2 to 5 percent in most Western countries, and it is rapidly increasing in Asiatic countries due to changes in dietary habits during the last years (1). Hyperglycemia is generally regarded as one of the major causes of vascular complications in diabetic patients. The endothelium is not a single-cell lining covering the internal surface of blood vessels but in fact plays a crucial role in regulating vascular tone and structure. Importantly, a healthy endothelium inhibits platelet and leukocyte adhesion to the vascular surface and maintains a balance of profibrinolytic and prothrombotic activity (2). Endothelial dysfunction has received increasing attention as a potential contributor to the pathogenesis of vascular disease in diabetes mellitus.

It has been reported that high extra-cellular glucose induces reactive oxygen species (ROS) production (2). Several studies have demonstrated that nitric oxide (NO) is abnormal in patients with type 2 diabetes (3). A multitude of experimental arguments have led to the concept that NO is not only involved in the control of vasomotor tone but also in vascular homeostasis function. NO is an endothelium-derived relaxing factor that produced through the conversion of arginin to citrulline by endothelial nitric oxide synthase (eNOS) or inducible nitric oxide synthase (iNOS) in vascular endothelial cells (4). The promoter region of eNOS gene contains tentative regulatory sequences, including cAMP, phorbol esters, and sterol-responsive elements (5). It has been shown that high-glucose concentration increases eNOS expression and NO generation in cultured human aortic endothelial cells (6). Increasing of NO production rates, often coupled with accelerated NO removal through poorly understood pathways, leads to impaired NO signaling and secondary generation of toxic NO-derived species (7). The reaction of NO with superoxide anion (O_2^-), yielding peroxynitrite ($ONOO^-$), accounts for a major part of the accelerated NO removal (7). Thus this reaction can have been an important role in enhanced rates of NO consumption as endothelium-derived relaxing factor.

In vitro studies have shown that various antioxidants, including vitamin E (α -tocopherol) and vitamin C (ascorbic acid), can prevent hyperglycemia-induced endothelial injury of dysfunction (8),(9). In addition, the epidemiological studies have demonstrated acute administration of vitamins improves endothelium-dependent vasodilatation (9), and there is an inverse relation between vitamin E and cardiovascular disease in diabetes patients (10).

Although a lot of studies have been done for protective role of antioxidants, especially vitamins, in the pathophysiology of diabetes and cardiovascular disease (9), (10), (11), (12), (13), (14), (15). but the literature data concerning the molecular mechanism of α -tocopherol and ascorbic acid effect on the production of NO are limited. We have previously shown that α -tocopherol and ascorbic acid are able to decrease LDL glycation as dose dependent in a high-glucose *in vitro* model (16). Therefore, our objective in this study is to evaluate of the effect of α -tocopherol and ascorbic acid on concentration of NOS, and eNOS and iNOS expression in high glucose-exposed human umbilical vein endothelial cells (HUVECs).

MATERIALS AND METHODS

Materials

Human umbilical vascular endothelial cells (HUVECs) were obtained from National Cell Bank Iran (NCBI), Pasteur Institute of Iran (Tehran, Iran). They were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 0.1 mg/ml heparin (Gibco BRL), 100 U/ml penicillin, 100 μ g/ml streptomycin, and 15% fetal bovine serum (Gibco BRL). The cultures were maintained at 37°C in an atmosphere of 95% air and 5% CO_2 . Medium was refreshed every three days. HUVECs of passages 3-5 were used for experiments. α -tocopherol, and ascorbic acid were purchased from Sigma (St. Louis, Mo, U.S.A), Dimethyl sulfoxide (DMSO) were obtained from Merck (Darmstadt, Germany).

Methods

Cell treatment and cell viability. In this study, HUVECs were treated with a medium containing 5.5 mM glucose (control) or 25 mM glucose (high glucose) for 24 hours (17), (18) in the presence or absence 50, 100, or 200 μ M of ascorbic acid or α -tocopherol. The incubation medium was then removed for analysis on NOS expression and NOS concentration, as described below. Cells viability were measured by adding 15 μ l of 2 mg/ml MTT (3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium) to cells in 96-well plates and incubating for 24 hour at 37°C. Then the medium was removed, 250 μ l DMSO was added to each well and the cells were incubated at 37°C for 4 hours. Absorption at 550 nm was read using a microplate reader (19).

Quantitative real-time PCR (RT-PCR) analysis. Total RNA was purified from HUVECs using the Qiagen RNA extraction kit (Germany). The RNA pellet was resuspended in RNase-free water and its concentration was determinate in a Lambda 18 Spectrometer (Perkin Elmer, Germany) at 260 nm. RNA (5 μ l) with concentration of 1 μ g was mixed with 2 μ l of dNTP, 2 μ l of 10 \times reverse transcriptase (RT) buffer, 2 μ l of random hexamer primer (Cinnagen, Iran), 1 μ l of ominiscript reverse transcriptase, and 8 μ l of RNase-free water. The RT mixture was incubated at 37°C for 60 minutes (method was done according to Qiagen cDNA kit, Germany made). The resulting cDNA was amplified by RT-PCR (ABI, USA) with the conditions of 95°C for 15 minutes (activation enzyme) and 45 cycles of 94°C for 15 seconds (denaturation), 55°C for 30 seconds (annealing), and 72°C for 34 seconds (extension). The primers were purchased from Qiagen Company (Germany). The primers for eNOS were 5'- GTG GCT GTC TGC ATG GAC CT-3' (upstream) and 5'- CCA CGA TGG TGA CTT TGG CT-3' (downstream), for iNOS were 5'-CGCCAAGAACGTGTTACCA-3' (upstream) and 5'-AGC AGG CAC ACG CCA TGA T-3' (downstream), and for β -actin, used as internal control, were 5'- AGC CTC GCC TTT GCCGA-3' (upstream) and 5'- CTG GTG CCT GGG GCG-3' (downstream). The PCR products were confirmed using melting-curve analysis for each gene. For comparison of transcript amounts between samples, a standard curve of cycle thresholds for serial

dilutions of a cDNA sample was established and then used to calculate relative levels of each gene [(20). All RT-PCR assays were performed in triplicate.

Assay of nitric oxide synthase (NOS) concentration. The concentration of NOS in cell culture was measured by enzyme-linked immunosorbent assay with commercial ELISA kits from Abnova Corporation (Taiwan). The cells that were used to measurement of NOS concentration were washed twice with Dulbecco's PBS, and cells were lysed with 400 μ l of lysis buffer (25 mM Tris/HCl (pH 7.6), 150 mM NaCl, 1 mg/l aprotinin, 10 mg/l leupeptin, 1 mM EDTA, 50 mM NaF, and 1% Triton-X-100). All samples were assayed in triplicate and expressed as pg/ml.

Statistical analysis

All data are shown as mean \pm SD. Treatment effects were analyzed using one-way analysis of variance (ANOVA). The level of significance was $p < 0.05$.

RESULTS

To verify inducement of high glucose on nitric oxide synthase (NOS), HUVECs were stimulated by 25 mM glucose concentration for 24 hours. High glucose increased the expression of eNOS mRNA and iNOS mRNA in comparison with low glucose (5.5 mM) approximately 92% and 78%, respectively (Figure 1). For determination of the best concentration of vitamins, we assayed the effect different concentrations (50 -200 μ M) of ascorbic acid and α -tocopherol on glucose (25 mM)-induced eNOS and iNOS expression for 24 hours. Ascorbic acid inhibited glucose-induced eNOS and iNOS mRNA expression in a significant dose-dependent manner ($p < 0.05$) when the concentration was limited between 50 and 100 μ M, whereas α -tocopherol showed a significant inhibition of glucose-induced eNOS and iNOS mRNA expression as dose dependent in concentrations of 50 to 200 μ M (Figure 2A,C). According to this study 100 μ M concentration of ascorbic acid is able to reduce eNOS and iNOS mRNA expression approximately 74% and 52%, respectively, but 200 μ M concentration of α -tocopherol is able to reduce eNOS and iNOS mRNA expression approximately 59% and 49%, respectively (Figure 2B,D).




Figure 1. The effect of 5.5 mM glucose concentration, as basal concentration, (black bar) and 25 mM glucose concentration (grey bar) on eNOS and iNOS expression assessed by RT-PCR in HUVECs after 24 hours incubation (37°C, 5% CO₂)
Data are represented as the mean \pm SD of triplicate determination. * $p < 0.05$ compared with control (black bar).

We investigated NOS concentration of HUVECs treated with 5.5 mM (low glucose) and 25 mM (high glucose) glucose (Figure 3). As shown in this Figure, NOS concentration in presence of high glucose as significant ($p < 0.05$), increase by 86% in comparison to the control (low glucose concentration). To assess whether ascorbic acid and α -tocopherol can inhibit high glucose-induced concentration of NOS, we examined NOS protein concentration by ELISA analysis. Treatment of cells (HUVECs) with ascorbic acid or α -tocopherol reduced the concentration of NOS induced by high glucose in a significant dose dependent manner ($p < 0.05$) (Figure 4A). This study showed that concentration of 100 μ M of ascorbic acid and 200 μ M of α -tocopherol are able to decrease concentration of NOS protein by 73% and 63%, respectively (Figure 4B).




Figure 2. The effect of 50, 100, and 200 μM concentrations of ascorbic acid (◆) and α-tocopherol (■) on eNOS (A), and iNOS (C) expression in presence high glucose concentration (25 mM); The comparison of inhibition percent of eNOS (B), and iNOS (D) expression in presence 25 mM glucose concentration and different concentrations (50-200 μM) of ascorbic acid and α-tocopherol. HUVECs treatment with vitamins at 37°C with air of 5% CO₂ for 24 hours

Data are represented as the mean±SD of triplicate determination. * $p < 0.05$ compared with control (in the absence of vitamins).




Figure 3. Effect of low glucose (5.5 mM), as basal concentration, and high glucose (25 mM) on concentration of NOS in HUVECs after 24 hours incubation (37°C, 5% CO₂)

Data are represented as the mean±SD of triplicate determination. * $p < 0.05$ compared with control (in the presence of low glucose).

Finally, according to linear (Pearson) analysis, correlation between eNOS and iNOS mRNA expression and NOS concentration in high glucose-induced HUVECs treated with ascorbic acid and/or α-tocopherol have been shown in Figure 5. This Figure, shows a significant positive correlation between concentration of NOS and expression of eNOS mRNA in presence ascorbic acid ($r = 0.99$; $p < 0.007$) (Figure 5A) and/or α-tocopherol ($r = 0.99$; $p < 0.007$) (Figure 5B). Additionally, a significant positive relation was also found between NOS concentration and iNOS mRNA expression with ascorbic acid ($r = 0.98$; $p < 0.02$) (Figure 5A) and/or α-tocopherol ($r = 0.99$; $p < 0.007$) (Figure 5B).




Figure 4. (A) The effect of 50, 100, and 200 µM concentrations of ascorbic acid (◆) and α-tocopherol (■) on HUVECs nitric oxide synthase (NOS) concentration in presence high glucose concentration (25 mM); **(B)** The comparison of inhibition percent of NOS protein concentration in presence high level of glucose (25 mM) and different concentrations (50-200 µM) of ascorbic acid and α-tocopherol. HUVECs treatment with vitamins at 37°C with air of 5% CO₂ for 24 hours

Data are represented as the mean±SD of triplicate determination. **p* < 0.05 compared with control (in the absence of vitamins)




Figure 5. Correlation between expression and concentration of eNOS (◆) and iNOS (■) in presence of high glucose (25 mM) and different concentrations (50 – 200 µM) of ascorbic acid (A) and α-tocopherol (B) in HUVECs

DISCUSSION

Dysfunction of endothelial cells are early manifestation of atherosclerosis that the most important result it is cardiovascular disease (21). Diabetes mellitus is an important risk factor for cardiovascular disease (2),(22). Pathways of nitric oxide (NO) signaling and reactions of lipid oxidation are important in both the maintenance of vascular homeostasis and the progression of vascular disease. Studies biochemical and cell biology have demonstrated that NO with oxidizing lipids could lead to either vascular protection or vascular injury (23), (24). The effect of NO is dependent to amounts it, if NO generated at low levels by nitric oxide synthase (NOS) can terminate propagating lipid radicals, reaction that would be protective. However, if NO generated at elevated levels, it can be converted to prooxidant species that can potentiate injury to vascular cells (24).

In this investigation, we showed that glucose could induce the expression of eNOS and iNOS mRNA in human vascular endothelial cells. Glucose (25 mM) increased the expression of eNOS 1.92-fold and iNOS 1.78-fold compare with control (5.5 mM glucose). These results are similar to previous studies that were reported by Cosentino et al. (6), Ding et al. (25), Zhu et al. (26), and Srinivasan et al.(27). In addition, according to Tesfamariam et al. report (28) it has been shown that concentration of high glucose is able by activating protein kinase C increase eNOS and iNOS genes expression in endothelial cells. According to results of this study and other investigations, we suggest that hyperglycemia may via increase of eNOS and iNOS gene expression in human vascular endothelial cells, accelerate atherosclerosis in diabetic patients. Several studies demonstrated that NO-derived reactive species, for example, peroxynitrite (ONOO⁻), nitrogen oxide (NO₂), and nitryl chloride (NO₂Cl) have an important role in

lipid oxidation and endothelial cells dysfunction (29), (30). Peroxynitrite is unique as a lipid oxidant, because it mediates oxidation of unsaturated fatty acids in the absence of transition metal catalysts (29). This NO-derived reactive species is more than two orders of magnitude potent than hydrogen peroxide in catalyzing lipid oxidation *in vitro* (31). Nitrogen oxide will both oxidize and nitrate unsaturated lipids (30), (32). These reactions result in formation of a complex mixture of products including nitrated lipid derivatives and alkyl nitrites. The oxidation of nitrite will yield nitryl chloride, which initiates lipid oxidation and can yield on LDL particle similar to that found in foam cells and so may be operative in atherogenesis (33), (34).

Vaziri *et al.* (35) demonstrated that antioxidant therapy ameliorated hypertension in rats, as well as, prospective studies have been shown cardiovascular benefits from antioxidant therapy (8), (9), (10). In the present study, ascorbic acid and α -tocopherol were used as protective agents against high glucose-induced NOS expression. It has been shown in several studies that antioxidants reduced vascular oxidative stress (36), (37). Additionally, ascorbic acid increased vasodilatation of forearm resistance arteries in humans with hypercholesterolemia (38), long-term smokers (39), essential hypertension (40), and coronary artery disease (41). Although numerous studies have shown antioxidants play a key role in oxidative stress inhibition. However, the literature concerning the effect vitamins on the expression and concentration of NOS in presence of high glucose is limited. In this investigation, we indicated that treatment of HUVECs with ascorbic acid and α -tocopherol decreased high glucose-induced eNOS and iNOS expression.

A dose-dependent manner was observed when the concentration of ascorbic acid and/or α -tocopherol was limited to the range of 50 to 100 μ M or 50 to 200 μ M, respectively. According to this study, optimum concentration for ascorbic acid and α -tocopherol obtained 100 μ M and 200 μ M, respectively. In presence of ascorbic acid (100 μ M) high glucose-induced eNOS and iNOS expression were inhibited by 74% (1.92-fold decreased to 1.1-fold) and 52% (1.78-fold decreased to 1.17-fold), respectively. α -tocopherol (200 μ M) also inhibited the expression of eNOS and iNOS in presence of high glucose by 59% (1.92-fold decreased to 1.21-fold) and 49% (1.78-fold decreased to 1.22-fold), respectively. According to these results, we suggest that 100 μ M ascorbic acid and 200 μ M α -tocopherol could restore eNOS and iNOS expression to levels observed in 5.5 mM glucose.

Our results are in agreement with the finding of several studies. For instance, Kang *et al.* (42) demonstrated that α -lipoic acid is able to decrease eNOS expression in bladder of rats with streptozotocin-induced diabetic. Similarly, Askwith *et al.* (43) also reported that *in vitro* taurine treatment reduces expression of NOS in high glucose-exposed human Schwann cells. In addition, there are other numerous studies that have been showed antioxidants, for example, melatonin, vitamin C (44), hesperidin (45), and vitamin E (46) could decrease expression of NOS isoforms *in vitro* and/or *in vivo*.

The mechanism of ascorbic acid and α -tocopherol on NOS downregulation is still unknown. However, according to study of Askwith *et al.* (43) the effect of antioxidants on NOS expression may be mediated by an antioxidant action, as well as, they also showed which this effect could be mediated by carbonyl scavenging or perhaps by restoring Ca^{+2} signaling (43).

In the present study, we also indicated that high glucose (25 mM) increase concentration of NOS protein in HUVECs by 86%, and found that ascorbic acid and α -tocopherol could reduce high glucose-induced NOS concentration in a significant and dose-dependent manner. The effective concentration of ascorbic acid and α -tocopherol that obtained in the present experimental study was 100 μ M and 200 μ M, respectively. These results confirmed our previous data in this study.

Finally, this investigation showed a positive and significant correlation between NOS protein content with NOS mRNA expression in HUVECs at present high glucose and ascorbic acid and α -tocopherol. According to these results, we suggest that ascorbic acid and α -tocopherol could decrease concentration of high glucose-induced NOS protein in HUVECs via NOS gene downregulation. Therefore, this is may be one of molecular mechanisms that ascorbic acid and α -tocopherol could prevent the increase in atherosclerosis induced by diabetes in human.

In conclusion, the results obtained in the present study show that both ascorbic acid and α -tocopherol reduced high glucose-induced expression of eNOS and iNOS in human endothelial cells. These vitamins appear to play an important role in the regulation of NOS with a mechanism of action which is at least partially independent of an antioxidant effect. These findings suggest molecular mechanism of these vitamins that possible preventive intervention for diabetes-accelerated atherosclerosis. Of course, further studies on regulation of ascorbic acid and α -tocopherol to signaling pathway is necessary.

Acknowledgements

This work was supported by a research grand from the Cellular and Molecular Research Center of Jundishapur University of Medical Sciences, Ahvaz, Iran (Project No. CMRC-59).

REFERENCES

- [1] Contreras F, Rivera M, Vasquez J, De la Parte M, Velasco M. Diabetes and hypertension physiopathology and therapeutics. *Journal of human hypertension*. **2000**;14:S26-S31.
- [2] Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. **2002**;105(9):1135-43.
- [3] Williams SB, Cusco JA, Roddy M-A, Johnstone MT, Creager MA. Impaired nitric oxide-mediated vasodilation in patients with non-insulin-dependent diabetes mellitus. *Journal of the American College of Cardiology*. **1996**;27(3):567-74.
- [4] Lamas S, Marsden PA, Li GK, Tempst P, Michel T. Endothelial nitric oxide synthase: molecular cloning and characterization of a distinct constitutive enzyme isoform. *Proceedings of the National Academy of Sciences*. **1992**;89(14):6348-52.
- [5] Marsden PA, Heng H, Scherer S, Stewart R, Hall A, Shi X, et al. Structure and chromosomal localization of the human constitutive endothelial nitric oxide synthase gene. *Journal of biological chemistry*. **1993**;268(23):17478-88.
- [6] Cosentino F, Hishikawa K, Katusic ZS, Lüscher TF. High glucose increases nitric oxide synthase expression and superoxide anion generation in human aortic endothelial cells. *Circulation*. **1997**;96(1):25-8.
- [7] Di Wang H, Hope S, Du Y, Quinn MT, Cayatte A, Pagano PJ, et al. Paracrine role of adventitial superoxide anion in mediating spontaneous tone of the isolated rat aorta in angiotensin II-induced hypertension. *Hypertension*. **1999**;33(5):1225-32.
- [8] Du X, Stockklauser-Färber K, Rösen P. Generation of reactive oxygen intermediates, activation of NF- κ B, and induction of apoptosis in human endothelial cells by glucose: role of nitric oxide synthase? *Free Radical Biology and Medicine*. **1999**;27(7):752-63.
- [9] Cummings PM, Giddens K, Nassar BA. Oral glucose loading acutely attenuates endothelium-dependent vasodilation in healthy adults without diabetes: an effect prevented by vitamins C and E. *Journal of the American College of Cardiology*. **2000**;36(7):2185-91.
- [10] Skyrme-Jones RAP, O'Brien RC, Berry KL, Meredith IT. Vitamin E supplementation improves endothelial function in type I diabetes mellitus: a randomized, placebo-controlled study. *Journal of the American College of Cardiology*. **2000**;36(1):94-102.
- [11] Golbidi S, Badran M, Laher I. Antioxidant and anti-inflammatory effects of exercise in diabetic patients. *Experimental diabetes research*. **2011**;2012.
- [12] Montonen J, Knekt P, Järvinen R, Reunanen A. Dietary antioxidant intake and risk of type 2 diabetes. *Diabetes Care*. **2004**;27(2):362-6.
- [13] Devasagayam T, Tilak J, Bloor K, Sane KS, Ghaskadbi SS, Lele R. Free radicals and antioxidants in human health: current status and future prospects. *Japi*. **2004**;52(794804):4.
- [14] Laight D, Carrier M, Ånggård E. Antioxidants, diabetes and endothelial dysfunction. *Cardiovascular research*. **2000**;47(3):457-64.
- [15] Ruhe RC, McDonald RB. Use of antioxidant nutrients in the prevention and treatment of type 2 diabetes. *Journal of the American College of Nutrition*. **2001**;20(sup5):363S-9S.
- [16] Ghaffari MA, Mojab S. In Vitro Effect of α -Tocopherol, Ascorbic Acid and Lycopene on Low Density Lipoprotein Glycation. *Iranian Journal of Pharmaceutical Research*. **2010**:265-71.
- [17] Ho FM, Lin WW, Chen BC, Chao CM, Yang C-R, Lin LY, et al. High glucose-induced apoptosis in human vascular endothelial cells is mediated through NF- κ B and c-Jun NH 2-terminal kinase pathway and prevented by PI3K/Akt/eNOS pathway. *Cellular signalling*. **2006**;18(3):391-9.
- [18] Ho FM, Liu SH, Liau CS, Huang J, Lin-Shiau SY. High glucose-induced apoptosis in human endothelial cells is mediated by sequential activations of c-Jun NH2-terminal kinase and caspase-3. *Circulation*. **2000**;101(22):2618-24.
- [19] Chen Y-J, Hsu K-W, Chen Y-L. Acute glucose overload potentiates nitric oxide production in lipopolysaccharide-stimulated macrophages: the role of purinergic receptor activation. *Cell biology international*. **2006**;30(10):817-22.
- [20] Turpaev K, Bouton C, Diet A, Glatigny A, Drapier J-C. Analysis of differentially expressed genes in nitric oxide-exposed human monocytic cells. *Free Radical Biology and Medicine*. **2005**;38(10):1392-400.
- [21] Vita J. Endothelial function and clinical outcome. *Heart*. **2005**;91(10):1278-9.

- [22]Haffner SM, Lehto S, Rönnemaa T, Pyörälä K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *New England journal of medicine*. **1998**;339(4):229-34.
- [23]Dusting GJ, Macdonald PS. Endogenous nitric oxide in cardiovascular disease and transplantation. *Annals of medicine*. **1995**;27(3):395-406.
- [24]Eiserich JP, Patel RP, O'Donnell VB. Pathophysiology of nitric oxide and related species: free radical reactions and modification of biomolecules. *Molecular aspects of medicine*. **1998**;19(4):221-357.
- [25]Ding Q, Hayashi T, Packiasamy AJ, Miyazaki A, Fukatsu A, Shiraishi H, et al. The effect of high glucose on NO and O²⁻ through endothelial GTPCH1 and NADPH oxidase. *Life sciences*. **2004**;75(26):3185-94.
- [26]Zhu M, Chen J, Tan Z, Wang J. Propofol Protects Against High Glucose-Induced Endothelial Dysfunction in Human Umbilical Vein Endothelial Cells. *Anesthesia & Analgesia*. **2012**;114(2):303-9.
- [27]Srinivasan S, Hatley M, Bolick D, Palmer L, Edelstein D, Brownlee M, et al. Hyperglycaemia-induced superoxide production decreases eNOS expression via AP-1 activation in aortic endothelial cells. *Diabetologia*. **2004**;47(10):1727-34.
- [28]Tefsamariam B, Brown ML, Cohen RA. Elevated glucose impairs endothelium-dependent relaxation by activating protein kinase C. *Journal of Clinical Investigation*. **1991**;87(5):1643.
- [29]Radi R, Beckman JS, Bush KM, Freeman BA. Peroxynitrite-induced membrane lipid peroxidation: the cytotoxic potential of superoxide and nitric oxide. *Archives of biochemistry and biophysics*. **1991**;288(2):481-7.
- [30]Gallon AA, Pryor WA. The reaction of low levels of nitrogen dioxide with methyl linoleate in the presence and absence of oxygen. *Lipids*. **1994**;29(3):171-6.
- [31]Patel R, Diczfalussy U, Dzeletovic S, Wilson M, Darley-Usmar V. Formation of oxysterols during oxidation of low density lipoprotein by peroxynitrite, myoglobin, and copper. *Journal of lipid research*. **1996**;37(11):2361-71.
- [32]O'Donnell VB, Eiserich JP, Chumley PH, Jablonsky MJ, Krishna NR, Kirk M, et al. Nitration of unsaturated fatty acids by nitric oxide-derived reactive nitrogen species peroxynitrite, nitrous acid, nitrogen dioxide, and nitronium ion. *Chemical research in toxicology*. **1999**;12(1):83-92.
- [33]Eiserich JP, Cross CE, Jones AD, Halliwell B, van der Vliet A. Formation of nitrating and chlorinating species by reaction of nitrite with hypochlorous acid a novel mechanism for nitric oxide-mediated protein modification. *Journal of Biological Chemistry*. **1996**;271(32):19199-208.
- [34]Hazell L, Arnold L, Flowers D, Waeg G, Malle E, Stocker R. Presence of hypochlorite-modified proteins in human atherosclerotic lesions. *Journal of Clinical Investigation*. **1996**;97(6):1535.
- [35]Vaziri ND, Ni Z, Oveisi F, Trnavsky-Hobbs DL. Effect of antioxidant therapy on blood pressure and NO synthase expression in hypertensive rats. *Hypertension*. **2000**;36(6):957-64.
- [36]Bauersachs J, Fleming I, Fraccarollo D, Busse R, Ertl G. Prevention of endothelial dysfunction in heart failure by vitamin E. *Cardiovascular research*. **2001**;51(2):344-50.
- [37]Chen X, Touyz RM, Park JB, Schiffrin EL. Antioxidant effects of vitamins C and E are associated with altered activation of vascular NADPH oxidase and superoxide dismutase in stroke-prone SHR. *Hypertension*. **2001**;38(3):606-11.
- [38]Ting HH, Timimi FK, Haley EA, Roddy M-A, Ganz P, Creager MA. Vitamin C improves endothelium-dependent vasodilation in forearm resistance vessels of humans with hypercholesterolemia. *Circulation*. **1997**;95(12):2617-22.
- [39]Heitzer T, Mu T. Antioxidant vitamin C improves endothelial dysfunction in chronic smokers. *Circulation*. **1996**;94(1):6-9.
- [40]Taddei S, Virdis A, Ghiadoni L, Magagna A, Salvetti A. Vitamin C improves endothelium-dependent vasodilation by restoring nitric oxide activity in essential hypertension. *Circulation*. **1998**;97(22):2222-9.
- [41]Gokce N, Keaney JF, Frei B, Holbrook M, Olesiak M, Zachariah BJ, et al. Long-term ascorbic acid administration reverses endothelial vasomotor dysfunction in patients with coronary artery disease. *Circulation*. **1999**;99(25):3234-40.
- [42]Kang DI, Kim SH, Lee SD, Kwak HS, Choi SH, Kim DR, et al. Effects of Alpha-lipoic acid on nitric oxide synthase expression and ultrastructural changes in the bladder of rats with streptozotocin-induced diabetes. *Korean Journal of Urology*. **2007**;48(2):212-8.
- [43]Askwith T, Zeng W, Eggo MC, Stevens MJ. Taurine reduces nitrosative stress and nitric oxide synthase expression in high glucose-exposed human Schwann cells. *Experimental neurology*. **2012**;233(1):154-62.
- [44]Sönmez M, Narin F, Akkuş D, Özdamar S. Effect of melatonin and vitamin C on expression of endothelial NOS in heart of chronic alcoholic rats. *Toxicology and industrial health*. **2009**;25(6):385-93.

- [45] Xiaoting L, Xiangyun Z, Shumei L, Minghua D, Liang X. Effect of hesperidin on expression of inducible nitric oxide synthase in cultured rabbit retinal pigment epithelial cells. *Retinal Degenerative Diseases: Springer*; **2010**. p. 193-201.
- [46] Calvisi DF, Ladu S, Hironaka K, Factor VM, Thorgeirsson SS. Vitamin E down-modulates iNOS and NADPH oxidase in c-Myc/TGF- α transgenic mouse model of liver cancer. *Journal of hepatology*. **2004**;41(5):815-22.