



Research Article

ISSN : 2277-3657  
CODEN(USA) : IJPRPM

## ***Antimicrobial Activity of Some Seaweed Collected from South East Coast of Jeddah, Saudi Arabia***

**Fotoon Sayegh<sup>1,2</sup>**

<sup>1</sup>Department of Biology, King Abdulaziz University, Jeddah 21589, Saudi Arabia

<sup>2</sup>Marine Natural Products Research Unit, King Fahd Medical Research Department, King Abdulaziz University, Jeddah 21589, Saudi Arabia

---

### **ABSTRACT**

Three marine seaweeds were collected from south east coast of Jeddah, Saudi Arabia. They were identified as species belonging to genera *Hypnea*, *Turbinaria* and *Padina*. They were extracted using hot water and five organic solvents. Their antimicrobial activities were determined against some clinical isolates of bacteria using disc diffusion assay. All the tested *Padina* and *Hypnea* extracts demonstrated activity against one or more of the tested bacteria. The methanolic extracts of *Padina* and *Hypnea* were the most active extracts compared to hot water and other organic extracts. They exhibited a broad spectrum of antibacterial activity against all the tested bacteria except *Mycobacterium* with MICs ranged from 125-400 µg/ml. Contrary to this, *Turbinaria* sp. did not show any detectable bactericidal activities. The toxicity for methanolic algal extracts was determined at the cell level by Biochemical Induction Assay and the Bioassay using *Artemia salina* as the test organism. No toxicity was detected against *A. salina* for the three tested algal extracts up to 300 µg/ml while at 400 µg/ml, the extracts of both *Turbinaria* sp. and *Hypnea* sp. showed some toxicity. By contrast, no toxicity for the methanolic algal extracts was detected by Biochemical Induction assay at 400 µg/ml. Only the extract of *Turbinaria* sp. had antitumor activity against Erlich cell line. The extracts showing good antimicrobial activities are undergoing further analysis to identify the active constituents which can be used as pharmaceutical compounds.

**Key words:** Seaweed, Jeddah Coast, Marine Algae, Antimicrobial Activity, Mic, Antitumor.

---

### **INTRODUCTION**

Seaweeds are renewable living resources used by common people as low calorie food, food additives, animal feed, and fertilizers in many parts of the world [1]. They have good nutritional importance because they are rich in carbohydrates, protein, vitamins, minerals and dietary fibers [2]. Because of increasing resistance of microorganisms to existing antibiotics, there is an increasing need for new antibiotics. Seaweeds or macro algae are considered as a source of secondary metabolites and bioactive compounds with antibacterial, antifungal, antiviral, antineoplastic, antifouling and anti-inflammatory activities. Green, brown and red algae offer particularly a rich source of natural and bioactive molecules [3-6]. They are providing valuable sources for the development of new drugs against cancer, microbial infections and inflammations [7-9].

Special attention has been reported for antiviral, antibacterial and/or antifungal activities related to marine algae against several pathogens [10]. As an efficient strategy of investigation, organic solvents have been used to extract the possible lipid-soluble active principles from macroalgae with activities against different human pathogens [11, 12]. Espechel et al. (1984) [13] indicated the antimicrobial activities of eight different marine algae collected in Southern Argentina. The extracts *Sargassum*, *Padina* and *Gracilaria* collected from different coastal regions of India suppressed the growth of some pathogenic bacteria and fungi [14]. Similarly, the methanolic extracts from thirty two

species of macroalgae from the Atlantic and Mediterranean coast of Morocco, belonging to Chlorophyta and Phaeophyta showed a significant capacity of antibacterial activities [15]. Since seaweeds offer particularly a rich source of bioactive molecules, they were screened for antimicrobial activities after extraction with organic solvents (Chloroform, ethanol, methanol and hexane).

Many tested algal extracts have many biological activities. Therefore, this study is made to screen three common algae, Padina, Hypnea and Turbinaria for their antimicrobial activities, toxicity, and antitumor productions. The most active isolated algal can be used in production of pharmaceutical compounds used in medicine.

## MATERIALS AND METHODS

### Seaweed sample collection

Algal samples were collected from Red Sea coast of Saudi Arabia at Shuaba, about 82 km south of Jeddah (longitude 35° 91' - 39° 24' East and latitude 15° 56' - 20° 52' North), during summer season, 2015 at depths ranging between 0.3 m and 1.5 m. They were collected in polyethylene bags by divers along the coast of the previous localities. Thereafter, they were transported wet to the lab in ice pack at 4°C. A complete sample of each collected algae was botanically identified at Biology Department, KAU, Saudi Arabia.

### Preparation of crude extracts

Each dried algal material (100 g) was powdered in a mortar and extracted two successive times at room temperature by continuous stirring for 16 h with 100 ml of different organic solvents including : acetone, chloroform, hexane, ethanol and methanol in addition to desterilized water. The organic extracts were collected and concentrated until dryness using rotary evaporator at 40°C but the water extract was lyophilized. The dried organic and water algal crude extracts were dissolved in 5 ml DMSO and screened for their biological activities.

### Microorganisms used

The antimicrobial activities of the different algal extracts were tested against different pathogenic and non pathogenic bacteria. They were obtained from culture collection of Laboratoire de Biochimie Microbienne, Fac. De Pharmacie, Nancy, France.

### The antimicrobial activities

The media used was nutrient agar for bacteria, Sabouraud dextrose agar for yeasts and fungi. The antimicrobial activity for each algal extract was detected against different test organisms. The used bacteria were Escherichia coli, Pseudomonas aeruginosa, Staphylococcus aureus, Klebsiella sp., Shigella sp., Corynebacterium michiganense and Mycobacterium. The technique used was agar well diffusion method described by Attaie et al., (1987) [16]. Petri dishes with 10 ml of nutrient agar were prepared, previously inoculated with 0.1 ml of a 24 h broth culture of tested bacteria. Four wells (6 mm) were made and each well was filled with 25 µl of the algal extract in DMSO. The inoculated plates were incubated for 24 h at 37°C. After incubation, the diameter of the inhibition zone was measured. Dimethyl sulfoxide (DMSO) was used as a negative control and Ampicillin as the standard antibacterial agent. The antimicrobial activity was taken on the basis of diameter of the zone of inhibition, and the mean of three readings was represented.

### The minimal inhibitory concentration

The minimal inhibitory concentration (MIC) was determined by the method suggested by Chand et al. (1994) [17] and modified by Aly and Gumgumjee (2011) [18] using 96 well ELISA trays. Each well was filled with 175 µl bacterial suspension,  $4 \times 10^6$  CFU ml<sup>-1</sup>, followed by 20 µl of the tested extracts in DMSO which was used as the control. The ELISA trays were incubated for 40 minutes, followed by addition of 5 µl of a 0.2% (w/v) Fluorescein diacetate in acetone and then the incubation was continued for 90 minutes more. The obtained green color was measured at 490 nm using MR7000 automatic ELISA tray reader.

### Antitumor activity of the algal extracts

Under a humidified atmosphere air with 5% CO<sub>2</sub>, Ehrlich carcinoma cell line was grown at 37°C for 48 h in RPMI 1640 medium (Sigma, USA) with 10 % fetal calf serum [18]. The collected cells were incubated with 400 µm/ml of the tested algal extract for 24 hrs., centrifuged for 2 min at 5000 rpm and counted using hemacytometer and trypan blue (Sigma, USA) in normal saline. LD<sub>50</sub> (lethal dose for 50 %) was determined and compared to the anticancer agent (Bleomycin).

### Toxicity of the algal extract

The toxicity of algal extracts was determined at the cell level using Biochemical Induction Assay, E. coli BR513 as test organism [19] and using Bioassay, Artemia salina as the test organism [20].

### Statistical analysis

Each experiment was carried out in three replicates, the mean of variable  $\pm$  standard deviation was calculated and significant differences between control and samples were calculated using t-test.

## RESULTS

Many investigations were carried out to discover algal products that inhibit pathogenic microorganisms including bacteria and fungi. The antimicrobial activities of three algal samples collected from Red Sea coast of Saudi Arabia at Shuaba, about 82 km south of Jeddah were investigated against some Gram positive and negative bacteria including *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella* sp., *Shigella* sp., *Corynebacterium michiganense* and *Mycobacterium* using agar well diffusion method. The tested algae were extracted using different solvents. The antimicrobial activity was measured by the zone of inhibition for all the different test organisms. The crude extracts of *Padina* and *Hypnea* revealed a wide range of antimicrobial activity against the tested pathogens while all the extracts of *Tepinaria* showed no antibacterial activities. The results were tabulated in the tables 1, 2 for the crude extracts of *Padina* and *Hypnea*, respectively.

The different extracts of *Padina* showed antibacterial activities against *Staphylococcus aureus* and *Escherichia coli*. The methanolic extract of *Padina* sp. Was the most active with antibacterial index of 80 (Table 1). The maximum inhibition was noticed against *E. coli* (15 mm) followed by *Pseudomonas aeruginosa* (13mm). Moderate antimicrobial activities were exhibited by ethanol, acetone, chloroform and hexane extracts. The lowest activities were obtained by water extract where the inhibition zone was ranged from 8-10mm and with antibacterial index of 50. None of the *Padina* extracts showed activities against *Mycobacterium* sp.

The methanolic extract of *Hypnea* showed promising antibacterial activities against all the tested bacteria especially *Escherichia coli*, *Pseudomonas* sp., *Shigella* sp., *Klebsiella* sp., *Bacillus subtilis* and *Staphylococcus aureus*. While chloroform and hexane extracts of *Hypnea* showed moderate antibacterial activities with inhibition zone ranged from 10-13mm. Weak antibacterial activity was obtained by water extract of *Hypnea* (diameter of the inhibition zone ranged from 8-9 mm). All *Hypnea* extracts showed no activity for *Mycobacterium*.

Minimal inhibitory concentration for the methanol extracts of the three tested algae were calculated for different test organisms (Table 3). The basic quantitative measures of the in vitro activity of the active substance are the minimum inhibitory concentration. The MIC is the lowest concentration of the active substance that results in inhibition of the visible growth under standard conditions. MIC of Ampicillin, calculated for bacteria, was ranged from 5-10  $\mu\text{g/ml}$ .

### Toxicity and antitumor activity

Methanol extracts of both *Turbinaria* and *Hypnea* showed a potent cytotoxic effect using standardised short-term toxicity test where *Artemia salina* was used as test organism. They showed mortality percentage  $>50\%$  (Table 4). The detected cell toxicity of *A. salina* was ranged from 9-29 % at 100-300  $\mu\text{g/ml}$  of the crude extracts. The obtained results showed that there was an increase in the toxicity of *Turbinaria* and *Hypnea* extracts with their elevated concentrations. The maximum toxicity of 65 and 55 % were observed at 400  $\mu\text{g/ml}$  of crude extract of *Turbinaria* and *Hypnea*, respectively. Cell toxicity was confirmed using Biochemical Induction Assay (BIA) and *E. coli* as test organisms. Both of the extracts could be a promising source of cytotoxic components. No toxicity was observed for the different concentration of methanol extract of *Padina* up to 400  $\mu\text{g/ml}$ .

No antitumor activity was recorded with the application of the crude extract of *Padina* and *Hypnea* species. The antitumor activity was observed for the methanolic extract of *Turbinaria* against Erlich ascites carcinoma cell line.

## DISCUSSION

Among marine organisms, algae are a large and diverse group of organisms from which a wide range of secondary metabolites have been isolated. A number of these compounds possess biological activity. Nagi et al., (2009) [21], Chiheb et al., (2009) [15] and Anbu-Jeba-Sunilson et al., (2009) [22] reported that marine algae are rich in active constituents showing antimicrobial activity. Furthermore, the ethanolic extract of *Sargassum ilicifolium*, *Padina tetrastomatica* and the methanolic extract of *Gracilaria corticata* showed maximum bacterial inhibition and the lowest antibacterial activities were noticed with chloroform extracts [14]. From Persian Gulf, the extracts of the collected red algae *Gracilaria salicornia* and *Hypnea flagelliformis* have strong antimicrobial activity due to the presence of acrylic acid, terpenes, heterocyclic compounds containing sulfur, chlorellin derivatives and halogenated aliphatic compounds [23]. Sastry et al. (1994) [24] noticed that algae have antibacterial activity against Gram-

positive and Gram-negative pathogenic strains after successive extraction with benzene, chloroform and methanol. Likewise, Mahasneh et al. (1995) [25] have shown antibiotic activity in organic extracts of six species of marine algae against multi-antibiotic resistant bacteria. Thus, it is evident from the results that these algae possess antimicrobial activity. Further studies are aimed at isolation and purification of phyto-constituents responsible for antimicrobial activity. Antimicrobial activity depends on algal type, the season of collection, and the efficiency of extraction of their active materials including used organic solvents and the method of extraction and assay.

Özdemir et al. (2001) [26] found that the *Spirulina* extracts, obtained from various solvents exhibited antimicrobial activity on both Gram-positive and Gram-negative organisms which may be indicative of the presence of broad spectrum antibiotic compounds. Numerous pharmacological active constituents like alkaloids, saponins, glycosides, tannins, chlorellin derivatives, acrylic acid and halogenated substances were identified as antimicrobial agents from algae. The results from the present study showed that Gram negative bacteria are more susceptible to algal extracts than Gram positive bacteria due to the differences in their cell wall structures and compositions, and this result was also supported from earlier works with different species of seaweeds [27]. Further in Gram-negative bacteria, the lipids of the outer membrane act as a target to many environmental substances including antibiotics, and the presence of a thick murine layer in the cell wall of Gram positive also prevents the entry of the inhibitors [28]. The overall antimicrobial activity assessed from the above results indicates the presence of active constituents in the extractions of seaweeds, thus they can be considered as potential sources of bioactive compounds acting as antibiotic or cytotoxic compounds.

In 21<sup>st</sup> century, cancer is a dangerous disease, it has increasing occurrence and it has caused about 25% of all the deaths in humans. Algal activity against cancer cell lines was documented, and many algae have shown cytotoxic and antitumor activities [29]. The methanolic extract of *Turbinaria* sp. acts as potential antitumor agents, while the extract of *Hypnea* and *Padina* has shown promising cytotoxic activity. Similarly, Barchi et al. (1984) [30] found out that lipophilic extracts of *Oscillatoria accutissima* were anti-neoplastic and showed toxic activities. *Codium tomentosum*, *Jania rubens* and *Padina pavonia* collected from Alexandria coast displayed 20% inhibition of tumor initiation [31]. The secondary products of algae, bromophenols, carotene and steroids showed antitumor activity [32]. Moreover, sulfated metabolites like fucoidans from the brown algae showed excellent activity against different tumor cell lines [33].

## CONCLUSION

This study reported the antimicrobial activity of different organic and water extracts of the 3 selected algae collected from Jeddah against some Gram positive and negative bacteria. Generally, the recorded antibacterial activity in the algal extracts was due to the active compounds that can be extracted, purified and used as pharmaceutical drugs. The algal extracts in different solvents exhibited different antimicrobial activities, and no solvent recorded effectively antibacterial activity against all the test bacteria.

## ACKNOWLEDGEMENTS

This project was funded by the Deanship of Scientific Research, King Abdulaziz University, Jeddah. The author thanks Dr. Naseem Radi, King Abdulaziz University, Faculty of Sciences, Biology Department for helping in algal collection and identification and Prof. Magda Aly, King Abdulaziz University, Faculty of Sciences, Biology Department for reading this manuscript.

## REFERENCES

1. Fuller, R., Roelofs, J. and Fryt, M. (1986). The importance of algae to stream invertebrates. *J. N. Am. Benthol. Soc.*, 5(4) :290-296
2. Rupérez P. and Saura-Calixto, F. 2001. Dietary fibre and physicochemical properties of edible Spanish seaweeds. *European Food Research and Technology* Volume 212, Number 3, 349-354.
3. Erickson, K. L. (1983). In : *Marine Natural Products* (edited by P. J. Scheuer), Academic Press, N.Y., 5, p. 131.
4. Bhakuni D.S and Rawat D.S. Ed. (2005). *Bioactive Metabolites of Marine Algae, Fungi and Bacteria*, in : *Bioactive Marine Natural Products*. Co-published by Springer 233 Spring Street, New York 10013, USA.

5. Katircioglu H., Beyatli Y., Aslim B., Yüksekdağ Z. and Atici T. (2006) Screening for Antimicrobial Agent Production of Some Microalgae in Freshwater. The Internet Journal of Microbiology, Volume 2, Number 2.
6. Hu Y., Chen J., Hu G., Yu J., Zhu X., Lin Y., Chen S., Yuan J. (2015). Statistical research on the bioactivity of new marine natural products discovered during the 28 years from 1985 to 2012. Mar. Drugs, 13 :202–221.
7. Prashantkumar, P Angadi SB, Vidyasagar GM (2006) Antimicrobial activity of blue-green and green algae Indian J. Pharmaceutical Science, Volume : 68 (5) 647-648
8. Bose, A., S. Sumanta, G. Jayanta Kumar, G. Tirtha, D. Gouri Kumar and S. Sudam, 2007. Analgesic, anti-inflammatory and antipyretic activities of the ethanolic extract and its fractions of *Cleome rutidosperma*. Fitoterapia, 78 : 515-520.
9. Blunt J.W., Munro M.H.G., Copp B.R., Keyzers R.A., Prinsep M.R. (2015). Marine natural products. Nat. Prod. Rep., 32 :116–211.
10. Singh, A., Thakur N.L. (2016) Significance of investigating allelopathic interactions of marine organisms in the discovery and development of cytotoxic compounds. Chem. Biol. Interact. ; 243 : 135–147.
11. Henriquez, P., Candia, A., Norambeuena, R., Silva, M. & Zemelman, R. (1979). Antimicrobial properties of marine algae. II. Screening of the Chilean coast for antimicrobial activity. Botanica mar., 22 : 451-453.
12. Shannon, E, Abu-Ghannam N. Antibacterial Derivatives of Marine Algae : An Overview of Pharmacological Mechanisms and Applications. Mar Drugs. 2016 Apr 22 ;14(4).
13. Espechel, M., Frailer, E.D. And Mayer, A.M. S. (1984). Screening of Argentine marine algae for antimicrobial activity. Hydrobiologia 116/117, 525-528 (1984).
14. Subba, Rangaiah G, Lakshmi P and Manjula E. (2010) Antimicrobial activity of seaweeds *Gracillaria*, *Padina* and *Sargassum* on clinical and Phytopathogens. International Journal of Chemical and Analytical Science, 1(6), 114-117.
15. Chiheb, I., Riadi H., Martinez L.J., Dominguez Selgar J.F., Gomez V.A., Bouziane H. and Kadiri M. (2009). Screening of antibacterial activity in marine green and brown macroalgae from the coast of Morocco. Afr. J. Biotechnol., 8 : 1258-1262.
16. Attaie, R., Whalen J., Shahani K.M., Amer, M.A., (1987). Inhibition of growth of *S. aureus* during production of acidophilus yogurt. J. Food Protec. 50, 224-228.
17. Chand S, Lusunzi I, Veal DAL, Williams R, Karuso P (1994) Rapid screening of the antimicrobial activity of extracts and natural products. J Antibiotics 47 :1295–1304
18. Aly MM, Gumgumjee N. (2011). Antimicrobial efficacy of *Rheum palmatum*, *Curcuma longa* and *Alpinia officinarum* extracts against some pathogenic microorganisms. Afr J Biotechnol ; 10 : 12058–63.
19. Elespuru RK, White RJ. (1983). Biochemical prophage induction assay : a rapid test for antitumor agents that interact with DNA. Cancer Res., 43(6) :2819–2830.
20. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlan JL (1982) Brine shrimp : a convenient general bioassay for active plant constituents. Plant Med 45 :31–34.
21. Nagi, A.A.H., I.M. Nurmas, N.S. Mariana, M. Habsah, S.V. Charles and S. Zambari, (2009). Antibacterial activity in Marine algae *Eucheuma denticulatum* against *Staphylococcus aureus* and *Streptococcus pyogenes*. Res. J. Biol. Sci., 4 : 519-524.
22. Anbu-Jeba-Sunilson, J., Suraj R., Anandarajagopal K., Rejitha G., Vignesh M. and Promwicht P. (2009). Preliminary phytochemical analysis, elemental determination and antibacterial screening of *Codium decorticatum*-a marine green algae. Int. J. Biol. Chem., 3 : 84-93.
23. Glombitza, K. W. (1979). Antibiotics from algae. In H. A. Hoppe, T. Levring & Y. Tanaka (eds), Marine Algae in Pharmaceutical Science. Walter de Gruyter, Berlin : 303-342.
24. Sastry, V.M.V.S. ; Rao, G.R.K. Antibacterial substances from marine algae : successive extraction using benzene, chloroform and methanol. Bot. Marina, 37 : 357-360, 1994.
25. Marasneh, I. ; Jamal, M. ; Kashasneh, M. ; Zibdeh, M. (1995). Antibiotic activity of marine algae against multi-antibiotic resistant bacteria. Microb., 83 : 23-26.
26. Özdemir, G., Conk Dolay, M., Küçükakyüz, K., Pazarbaşı, B., Yilmaz, M. (2001). Determining the antimicrobial activity capacity of various extracts of *Spirulina platensis* produced in Turkey's conditions. J Fisheries and Aquatic Sciences 1st. Algal Technology Symposium. 18/1, 161-166.

27. Tajbakhsh, S, Pouyan M, Zandi K, Bahramian P, Sartavi K, Fouladvand M, Asayesh G, Barazesh A. In vitro study of antibacterial activity of the alga *Sargassum oligocystum* from the Persian Gulf. *Eur Rev Med Pharmacol Sci.* 2011 Mar ;15(3) :293-8.
28. Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. *Cold Spring Harb Perspect Biol.* 2010 May ; 2(5) : a000414.
29. Zandi, K, Tajbakhsh, S., Nabipour, J., Rastian, Z., Yousefi, F., Sharafian S. and Sartavi K. (2010). In vitro antitumor activity of *Gracilaria corticata* (a red alga) against Jurkat and molt-4 human cancer cell lines. *African Journal of Biotechnology* Vol. 9(40), pp. 6787-6790.
30. Barchi, J.J., Moore, J.R.E., Patterson, G.M.L. (1984). Accutiphycin and 20, 21-Didehydroaccutiphycin, new antineoplastic agents from the cyanophyte *Oscillatoria acutissima*. *J Am Chem Soc.* 106, 8193-8197.
31. El-Masry, M. H., Mostafa, M. H., Ibrahim, M.A., El-Naggar, M. (1995) Marine algae that display anti-tumorigenic activity against *Agrobacterium tumefaciens*. *FEMS Microbiology Letters*, Volume 128, Issue 2, pages 151–155.
32. Xu, N, Fan X, Yan X, Tseng C (2004). Screening marine algae from China for their antitumor activities. *J. Appl. Phycol.* 16 : 451-456.
33. Ly BM, Buu NQ, Nhut ND, Thinh PD, Thi T, Van T (2005). Studies on fucoidan and its production from Vietnamese brown sea weeds. *AJSTD*, 22 : 371-380.

**Table 1.** Antimicrobial activity of *Padina* extracts obtained with different organic solvents against different tested bacteria

Tested microorganisms	Inhibition zones (mm) of the different algal extracts**					
	Methanol	Ethanol	Acetone	chloroform	Hexane	H <sub>2</sub> O (control)
<i>Escherichia coli</i>	19±0.1	10±0.1	11±0.2	10±0.1	10±0.0	8±0.6
<i>Pseudomonas aeruginosa</i>	15±0.2	13±0.3	10±0.1	10±0.1	11±0.2	8±0.2
<i>Shigella</i> sp.	13±0.3	12±0.2	10±0.1	15±0.2	11±0.1	8±0.1
<i>Klebsiella</i> sp.	11±0.2	12±0.1	10±0.2	10±0.1	11±0.1	8±0.1
<i>Bacillus subtilis</i>	11±0.1	11±0.1	11±0.1	10±0.3	10±0.0	8±0.0
<i>Staphylococcus aureus</i>	11±0.1	10±0.2	13±0.2	10±0.2	15±0.2	10±0.2
<i>Mycobacterium</i>	-	-	-	-	-	-
Antibacterial index***	80*	68*	65*	65*	68*	50

\*: significant results compared to control, \*\* Inhibition zone ,7–10-mm =weak activity ; 10–15-mm = good activity; (15–20 mm very good activity and (-) =No activity. \*\*\* Activity index was calculated as the sum of the net values of inhibition zone diameters (mm) against the tested bacterial strains

**Table 2.** Antimicrobial activity of different *Hypnea* extracts obtained with different organic solvents against different tested bacteria

Microorganisms	Inhibition zones (mm,) against investigated extracts**					
	Methanol	Ethanol	Acetone	chloroform	Hexane	H <sub>2</sub> O (control)
<i>Escherichia coli</i>	12±0.0	11±0.1	11±0.1	10±0.3	10±0.0	8±0.1
<i>Pseudomonas aeruginosa</i>	12±0.2	09±0.1	11±0.1	10±0.1	11±0.2	8±0.3
<i>Shigella</i> sp.	12±0.1	10±0.1	12±0.2	12±0.2	11±0.1	9±0.2
<i>Klebsiella</i> sp.	12±0.1	09±0.1	07±0.1	13±0.3	11±0.1	8±0.1
<i>Bacillus subtilis</i>	12±0.0	10±0.1	10±0.1	13±0.3	10±0.0	8±0.1
<i>Staphylococcus aureus</i>	12±0.1	10±0.1	10±0.0	10±0.3	15±0.2	8±0.2
<i>Mycobacterium</i> sp.	-	-	-	-	-	-
Antibacterial index***	74*	59*	61*	68*	68*	49

\*: significant results compared to control, \*\* Inhibition zone ,7–10-mm =weak activity ; 10–15-mm = good activity ; (15–20 mm very good activity and (-) =No activity. \*\*\* Activity index was calculated as the sum of the net values of inhibition zone diameters (mm) against the tested bacterial strains

**Table 3.** The minimal inhibitory concentrations of the methanolic algal extracts in addition to Ampicillin (control) for different tested bacteria

Minimal inhibitory concentration (MIC, µg/ml)				
Microorganisms	Control antibiotic (Ampicillin)	Padina sp.	Turbinaria sp.	Hypnea sp.
<i>Escherichia coli</i>	10	100	>400	250
<i>Pseudomonas aeruginosa</i>	5	125	>400	250
<i>Shigella</i> sp.	5	125	>400	250
<i>Klebsiella</i> sp.	5	150	>400	250
<i>Bacillus subtilis</i>	2	250	>400	200
<i>Staphylococcus aureus</i>	2	250	>400	250
<i>Mycobacterium</i>	>400	>400	>400	>400

**Table 4.** The toxicity and antitumor activity of the three methanolic algal extracts

Conc. (µg/ml) Algal extract	Toxicity (LD <sub>50</sub> , µg/ml)	Toxicity (Biochemical Induction Assay)	Antitumor Activity against Erlich ascites carcinoma (400 µg/ml)
Control (Bleomycin)	>400	-	+
<i>Padina</i> sp.	400>	-	-
<i>Turbinaria</i> sp.	375*	+	+
<i>Hypnea</i> sp.	350*	-	-