

## Evaluation of Counter Irritant Potential of Aqueous Bark Extract of *Cinnmon Loureiroi*

Imran Ahmad Khan<sup>1\*</sup> Abdul Aziz<sup>2</sup> Shaukat Hussain Munawar<sup>3</sup> Zahid Manzoor<sup>4</sup> Muzammal Sattar<sup>5</sup> Aqsa Afzal<sup>6</sup>

<sup>1,2,6</sup> Faculty of Pharmacy, Bahauddin Zakariya University, Multan, Pakistan

<sup>3,4</sup> Faculty of Medicine and Allied Medical sciences, Isra university, Islamabad, Pakistan

<sup>5</sup> Department of Physiology & Pharmacology, University of Agriculture, Faisal Abad, Pakistan  
imranahmadkhadurrani@gmailcom

Subject: Pharmacology

### Abstract

Crude aqueous bark extract of *Cinnamon loureiroi*. was evaluated for counter irritant activity. Irritation was induced by the clockwise frictional movement of fine sand paper to the ear of rabbits of average 1.5 kg body weight. The counter-irritant activity was determined by calculating the mean decrease in redness and erythma with those of control. cinnamon extract (25, 50, 75, 100µg / 10 ml) showed counter irritant when compared with standard drug dexamethasone. All the extracts, showed the counter irritant activity. Highest (91.97% inhibition) and the lowest (41.39 % inhibition) respectively.

**Keywords:** Counter irritant, *Cinnamon loureiroi* , rabbit's skin, dexamethasone .

### Introduction

Skin irritation is known as “the production of reversible damage of the skin following the application of a test substance for up to 4 hours” (OECD, 1981). Pathological characteristics and manifestation of skin irritation such as, Erythema and edema are manifestations of dermal irritation. Irritation is initially manifest by redness (erythema), vesicles, serous exudates, serous scabs (eschar) and various degrees of swelling (edema). Over time, other reactions may be manifest, like small areas of scaling, hyperplasia, hyperkeratosis and alopecia. Histopathology is useful in discerning among responses. In most cases inflammation is well developed within the first 72 hours of observation. ( Gallegos et al., 2006)

*Cinnamon loureiroi* use as medicine is thousand years old mentioned in several books of the Bible and in the histories of ancient Rome and Egypt as well as medieval Europe ( Keith Singletary, 2008). Ayurvedic and folklorik uses are, wound healing, flatulance, erectile sysfuction, conjectivitis, leukorrhea, vaginitis, rheumatism, neuralgia, aphrodisiac, anti allergy, antifungal, insecticidal, antipyretic, analgesic, antiulserant, nematocidal (www.naturalstandard.com; Das et al., 2013). Its historical uses were, antidiabetic ( Qin et al., 2003; Kim et al., 2006) anticancer (Schoene et al., 2005 )

antimicrobial ( Sing et al., 2007) anti inflammatory (Kim et al., 2007) blood pressure lowering ( Preuss et al., 2007) cholesterol lowering ( Khan et al., 2003) antidiarrheal, cough, sore throat, indigestion, , chest congestion, abdominal pain, headache (waris et al., 2003 ). tooth ache ( Archer, 1998) medication resistan yeast infections (www.herbwisdom.com). Smelling of cinnamon enhances cognitive function and memory ( Palmer et al., 1998). prevention of cardiovascular diseases, carcinogenesis, atherosclerosis (Srinivasan, 2005) vomiting (Khan et al.,2014)

### Material and Method

#### Material

##### Chemicals

Dexamethazone sod. Was perched from Ethical laboratories (Pvt) Ltd. Pakistan and *Cinnamon loureiroi* was purchased from local market of Multan, Pakistan.

##### Animals

Rabbits of either sex with the average weight of 1.5 kg was purchased from Animal market Hussain Agahi Multan, After 1 hr observing the nomal dermatological and allergic behaviour, the counter

irritant activity was evaluated. All the rabbits were kept under laboratory conditions at room temperature with 12h light and dark cycles. All animal experiments were carried out in accordance with the acts of the Animal Ethical Committee of Baha-uddin zakariya university, Multan, Pakistan.

## Method

### Collection of Plant Materials

Indigenous medicinal plant *Cinnamon loureiroi*. known by a local name of “Dal chinni”. The plant were collected from the local market of Multan, Pakistan. The plant material was authenticated by expert taxonomist, Professor Dr. Altaf Dasti at the Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan.

### Crude extract

The plant material was made free from foreign adulterants and vegetative debris by hand picking. Special electrical herbal Grinder was used to form coarse powder. Uniform dark brown powder was obtained with characteristic smell. Powdered *Cinnamon loureiroi* dissolved in distilled water, fluid obtained was filtered through Whatman-1 Filter paper (Horbom, 1971).

### Counter- irritancy assay

Assay for counter irritant effect of aqueous bark extract of *Cinnamon loureiroi*. was performed with some specified modifications in assay described by Syed Saeed ul Hassan et al. ( 2013). Sand paper with fine particles was used to irritate inner surface of rabbit's ear in clockwise direction for 10 minutes. Irritation, redness and erythema were produced in area of 2.0 cm<sup>2</sup> in diameter. 100µl solution from each concentration of *Cinnamon loureiroi* and dexamethason (Standard) was applied to the irritated area. Distilled water treated ear was used as control. Ears were examined for intensity of erythema. A group of 3 rabbits for each was used while performing main assay by increasing concentration of irritants. Rabbits were examined after every 10 minutes. The numbers of ear showing decreased irritancy, redness and erythema were observed and recorded. The previous authors used control and test extract but not the effect of standard drug comparison. The time, dose and degree of counter irritancy was missing which is provided in this assay.

### Phytochemical Study

The crude plant extracts were initially screened qualitatively with different organic solvents and

reagents to detect the presence of some phytochemicals classes (Tona et al., 1998).

### Toxicity Study

*Cinnamon* is used as a spice in food material in Asia so its safety is quite obvious. Budavari et al.(1989) have reported acute toxicity of *Cinnamon* in the animals is very low i.e. Benzaldehyde (LD50 orally, 1300 mg/kg rat), cinnamaldehyde (LD50 orally, 2220 mg/kg rat), linalool (LD50 orally, 2790 mg/kg rat), and salicylaldehyde (LD50 orally, 520 mg/kg rat)


## Statistical analysis

### Results


Preliminary phytochemical screening detected presence of tannins, phenols, saponins alkaloid, anthraquinones and coumarins as constituents of the crude aqueous bark extract of *Cinnamon loureiroi*. (Cl.Cr).

**Table 1: Pytochemical analysis of *Cinnamon loureiroi* (bark) crude extracts (Cl.Cr).**

Sr. no	Test	Observations	Result
1	Alkaloid	Ppt	Positive
2	Saponins	1cm froth	Positive
3	Tannins	Light purple	Positive
4	Anthraquinones	Pink	Positive
5	Coumarins	Yellow fluorescence	Positive
6	Phenols	Light purple	Positive
7	Flavanoid	Light yellow colour	Positive



**Figure. 1: Counter irritancy effect, Group-I: Control (Distilled water); Group-II: Standard drug (Dexamethasone); Group-III; . Cinnamon loureiroi (50, 100 µl)**



**Figure 2:** Counter irritancy time, Group-I: Control (Distilled water); Group-II: Standard drug (Dexamethasone); Group-III; Cinnamon loureiroi. .(50, 100  $\mu$ l )

**Table 2:** Counter irritant activity of aqueous bark extracts of *Cinnamon loureiroi*. S.E.M.= Standard Error of Mean, \* $P < 0.1$  and \*\* $P < 0.005$  vs. control showing significant and most significant values using unpaired Student's  $t$ -test

Groups	Mean degree of Erythma $\pm$ S.E.M	Inhibition (%) of Erythma	Countergerency time
Control 10ml/kg	91 $\pm$ 1.66		
Cinnamon 100 ug/10 ml	7.3 $\pm$ 3.12	91.97% ***	10min
Cinnamon 75 ug/10 ml	15.56 $\pm$ 1.21	82.90% ***	30min
Cinnamon 50 ug/10 ml	23 $\pm$ 1.89	74.72% **	50min
Cinnamon 25 ug/10 ml	53.33 $\pm$ 2.31	41.39% *	120 min
Dexamethason	1 $\pm$ 1.09	98.90% ***	1min



**A**



**B**



**C**



**D**



**E**



**F (Dexa)**

**Figure 3 (A-F): Counter irritant activity Cinnamon and Dexamethasone ( Standarded drug)**

## Discussion

Positive results are connected to the presence of tannins and saponins in the *Cinnamon*. On local application, the tannins act as astringents, healing, antiexudative, anti-irritative, anti-inflammatory antiseptic, anesthetic and antioxidant. Tannins form complexes with the proteins from the superficial layers of the skin, leading to the formation of a protective layer of protein-tannins. Moreover, the tannins act as antiseptics through the precipitation of the proteins from the membrane of the microorganisms and anti-inflammatory through the inhibition of the synthesis of prostaglandins and the freeing of the plachetary activating factor (PAF) (Mahesh et al., 2008). Saponins, which from a structural point of view are glycosides, have an antiseptic and antimicrobial action, in a non-harmful way for the neighboring cellular tissues. The specialty literature draws attention to. Flavonoic derivates develop anti-inflammatory effects (mainly by inhibiting the freeing of lisosomal enzymes and reducing the level of oxygen-reactive species) (Casadevall et al., 2001), anti-allergic effects (by inhibiting the classical way the seric complement is activated), anti-microbial, capillary-protective and antioxidant.

## Acknowledgement

Authors are thankful to Dr.Khaild Hussain Janbaz and Fatima Saqib for their expert opinion and guidance throughout the experiement.

## “Cite this article”

I.A. Khan, A. Aziz, S. H. Munawar, Z. Manzoor M. Sattar, A. Afzal “Evaluation of Counter Irritant Potential of Aqueos Bark Extract of Cinnmon Loureiroi”. Int. J. of Pharm. Res. & All. Sci, 2014;3(1), 30-35

## Reference

- (1) Ana Gallegos Saliner, Grace Patlewicz & Andrew P. Worth (2006). Review of Literature-Based Models for Skin and Eye Irritation and Corrosion .
- (2) Organisation for Economic Co-operation and Development (1981). Guideline for Testing of Chemicals No. 404: *Acute dermal irritation/corrosion*. OECD, Paris, France, adopted on 12 May 1981,
- (3) Keith S (2008). Cinnamon: Overview of Health Benefits . Nutr Today.43: 263–266
- (4) Natural Standard: The Authority on Integrative Medicine. [www.naturalstandard.com](http://www.naturalstandard.com)
- (5) Manosi S, Mandal S, Mallick B and Hazra J (2013). Ethnobotany, Phytochemical and pharmacological aspect of Cinnamomum Zeylanicum.int. Res. J. Pharm. 4: 58-63
- (6) Qin B, Nagasaki M, Ren M, Bajotto G, Oshida Y and Sato Y ( 2003). Cinnamon extract (traditional herb) potentiates in vivo insulin-regulated glucose utilization via enhancing insulin signaling in rats. Diabetes Res Clin Pract. 62: 139-148
- (7) Kim S, Hyun S and Choung S (2006). Anti-diabetic effect of cinnamon extract on blood glucose in db/db mice. J Ethnopharmacol. 104: 119-123.
- (8) Schoene N, Kelly M, Polansky M and Anderson R (2005). Water-soluble polyphenols from cinnamon inhibit proliferation and alter cell cycle distribution patterns of hematologic tumor cell lines. Cancer Lett. 230:134-140.
- (9) Singh G, Maurya S, Lampasona M and Catalan C (2007). A comparison of chemical, antioxidant and antimicrobial studies of cinnamon leaf and bark volatile oils, oleoresins and their constituents. Food Chem Toxicol. 45:1650-1661.
- (10) Kim D, Kim C, Kim M, et al. (2007). Suppression of age-related inflammatory NF-κB activation by cinnamaldehyde. Biogerontology. 8: 545-554
- (11) Preuss H, Echard B, Polansky M and Anderson R (2006). Whole cinnamon and aqueous extracts ameliorate sucrose-induced blood pressure elevations in spontaneously hypertensive rats. J Am Coll Nutr. 25: 144-150
- (12) Khan A, Safdar M, Ali Khan M, Khattak K and Anderson R (2003). Cinnamon improves glucose and lipids of people with type 2 diabetes. Diabetes Care. 26: 3215-3218
- (13) Waris Q, Salman R A, Raheem H D, Sana J, Aysha N and Ammara R (2003). Use of folk remedies a mong patients in Karachi pakistan. JAMC. 15-21
- (14) Archer AW (1988). Determination of cinnamaldehyde, coumarin and cinnamyl alcohol in cinnamon and cassia by high-performance liquid chromatography. J. Chromatogr. 447: 272–276.
- (15) <http://herbwisdom.com>
- (16) Palmer AS, Stewart J and Fyfe L (1998). Antimicrobial properties of plant essential oils

- and essences against five important food borne pathogens. Lett. Appl. Microbiol. 26: 118 – 122.
- (17) Srinivasan K ( 2005). Role of spices beyond food flavouring: nutraceuticals with multiple health effects. Food Rev. Int. 21: 167–188.
- (18) Harborne JB (1973). Methods of plant analysis. In, Phytochemical Methods. Chapman and Hall, London, 1-7.
- (19) Syed Saeed ul Hassan, Imran Waheed, Muhammad Khalil-ur-Rehman, Uzma Niaz and Muhammad Asif Saeed. Counter irritant activity of *Carthamus oxycantha* Pak. J. Pharm. Sci., Vol.26, No.4, July 2013, pp.665-672
- (20) Tona L, Kambu K, Ngimbi N, Cimanga K and Vlietink AJ (1998). Antiamoebic and Phytochemical Screening of Some Congolese Medicinal Plants. Planta Med. 61: 57-65.
- (21) Budavari SB, O'Neil MJ, Smith A and Heckelman PE (1989). The Merck Index. Merck and Co, Rahway, NJ.
- (22) Jayaprakasha GK, Raom LJ and Sakariah KK (2002). Chemical composition of volatile oil from *Cinnamomum zeylanicum* buds. Zeitschrift für Naturforschung C. Journal of Biosciences 57: 990–993.
- (23) Sandigawad AM and Patil CG (2011). Isolation and Characterization of Alkaloids from *Cinnamomum* Scha.(Lauraceae) species. Advances in Bioresearch. 2: 90 - 91
- (24) [http:// www.botanical-online.com](http://www.botanical-online.com)
-