



Research Article

ISSN : 2277-3657
CODEN(USA) : IJPRPM

Antibacterial Activity of Taxifolin Isolated from Acacia Catechu Leaf Extract– An in Vitro Study

Thanish Ahamed S.¹, Lakshmi T.²

¹ Graduate Student, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Chennai-600077, India

² Professor, Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, 162, Poonamallee High Road, Chennai-600077, India

ABSTRACT

Background : An anti-microbial is an agent which causes the death or hinders the growth of microorganisms such as bacteria and fungi. Acacia catechu that is also commonly called Mimosa catechu, is a deciduous, thorny tree which grows up to be 15 m (50 ft) high. In folk medicine, Acacia catechu is applied as a treatment of many illnesses and specially for mother and child healthcare. The main constituent of Acacia catechu is Taxifolin which contains antifungal, antiviral, antibacterial, anti-inflammatory and anti-oxidant property. Aim : The study aimed at evaluating the anti-microbial activity of taxifolin isolated from Acacia catechu leaf extract against Streptococcus mutans and Lactobacillus acidophilus, which may help in development of health products without artificial chemical agent.

Key words: Antimicrobial, Herbal, Antibacterial, Acacia, S. Mutans, L. Acidophilu.

INTRODUCTION

Medicinal plants have been used for centuries as a remedy for various human diseases [1] through having antibacterial, antifungal or antioxidant activities [2]. Due to the fact that bacteria have been increasingly developed resistance to the currently available antibiotics, the need to find new antibacterial agents has been raised [3, 4]. Anti-microbial activity of certain plants has a significant effect on various microorganisms.

An antimicrobial can be defined as an agent that causes the death of microorganisms or hinders their growth with the least damage to the host cells [5]. Anti-microbial agents play a major role in maintaining good health. The management of infections like pneumonia, tuberculosis, malaria, and AIDS has been threatened by antimicrobial resistance. So, it is important to develop stronger antimicrobial agents which are achieved by various anti-microbial agents. One such antimicrobial agent is taxifolin which is isolated from many trees such as Acacia catechu, conifers like the Siberian larch, Larix sibirica in Russia, and Chinese yew [6].

Acacia catechu commonly known as catechu is a medicinal plant used for varied purposes [7]. The bark of this plant is powerful antioxidant [8], astringent, anti-inflammatory, anti-bacterial and antifungal in nature [9]. Sore throats and diarrhoea, have been treated by the extract of this plant, and also in cases of high blood pressure, dysentery, colitis, gastric problems, bronchial asthma, cough, leucorrhoea and leprosy, it is considered to be useful [10, 11]. It has also been used as mouthwash for mouth, gum, sore throat, gingivitis, dental and oral infections. The current study has been done to evaluate the anti-microbial activity of taxifolin isolated from Acacia catechu against Streptococcus mutans and Lactobacillus acidophilus

Taxifolin is a flavanone, a type of flavonoid. They are isolated from many trees. One of those tree is Acacia catechu [12]. Taxifolin is an anti-cancer drug. By inhibiting the fatty acid synthase in cancer cells, they are able to prevent

the growth and spread of cancer cells. The property of taxifolin to activate fibril formation and improve the stabilization of fibrillar forms of collagen can be implemented in medicine.

Streptococcus mutans and Lactobacillus acidophilus are the most important microbes present in the oral cavity. They are one of the major reasons for the dental caries. They are gram positive bacteria [13, 14]. The lactobacilli and streptococci which are generally named the lactic acid bacteria are the most important genera of the group of bacteria, so it is not surprising that microbial species within these genera have been proposed as specific agents of the acid production that is primary to the dental caries process [15].

The study has been conducted to evaluate the anti-microbial activity of taxifolin extracted from Acacia catechu against Streptococcus mutans and Lactobacillus acidophilus

Drugs

Taxifolin was purchased from Sigma-Aldrich, India.

Acacia catechu – obtained from green chemical herbal extracts and formulation, Bengaluru.

Test microorganism

Streptococcus mutans - ATCC 25175 Lactobacillus acidophilus - ATCC 4356 from Himedia, Mumbai.

MATERIALS AND METHODS

Antibacterial activity

To screen their susceptibility, active cultures were prepared by transferring loop full of cells from stock cultures to test Mueller Hilton Broth containing tubes, and then, were incubated at 37°C for 24 hours [16, 17].

Disc diffusion method

Onto the plates containing sterile Mueller Hinton Agar, 0.1mL of the bacterial culture was swabbed uniformly. Different concentrations of the sample were loaded a day before on 5 mm sterile discs. These discs were placed on the bacteria inoculated plates. The plates were incubated at 37°C for 24 hours. After incubation period, the diameter of inhibition zones which was formed around the discs was measured in millimeter. The study was performed in duplicates for all the samples.

RESULTS

Taxifolin –Antibacterial activity

Table 1. Antibacterial activity of taxifolin against S. mutans and L. acidophilus

Concentration (mg/ml)	Streptococcus mutans			Lactobacillus acidophilus		
	Zone of inhibition (in mm)			Zone of inhibition (in mm)		
	Plate 1	Plate 2	Average	Plate 1	Plate 2	Average
1.5	18	16	17	7	7	7
2.0	19	17	18	7	8	7.5
2.5	22	24	23	13	16	14.5

Control

Table 2. Antibacterial activity of chlorhexidine

Concentration (mg/ml)	Chlorhexidine		
	Zone of inhibition (in mm)		
	Plate 1	Plate 2	Average
1.5	24	26	25
2.0	25	20	22.5
2.5	24	22	23

The antibacterial activity of the Taxifolin from Acacia catechu leaf was tested against S. mutans and L. acidophilus using the disc diffusion method. As shown in Table 1, three different concentrations of 1.5mg/ml ,2mg/ml ,2.5mg

/ml were used in this study. Taxifolin inhibits *Streptococcus* mutants significantly when compared to *Lactobacillus acidophilus*. Chlorhexidine is a gold standard drug for maintaining oral hygiene with its antibacterial effect. The Taxifolin showed moderate antibacterial efficacy against *S. mutans* & *L. acidophilus* when compared to the control chlorhexidine.

DISCUSSION

Taxifolin which is isolated from *Acacia catechu* has various medical properties. Various studies have been done to disclose the medical property of taxifolin. Taxifolin has been found as anti-cancer drug. The therapeutic promise of dihydroquercetin(taxifolin) in major inflammatory disease states such as cancer was recently reviewed by [18, 19]. Further studies have been done, and the results concluded that Taxifolin and possibly other flavonoids with a similar molecular structure may act as “enhancers” in combination with Andro to treat prostate cancer [20].

Taxifolin has been said to have a dose-dependent effect on inhibiting the ovarian cancer cells. It also has a strong correlation between the anti-proliferative effects of DHQ derivatives on murine skin fibroblasts and human breast cancer cells [21]. They have been used positively to treat patients infected with strains of methicillinresistant *S. aureus* (MRSA) [22]. They increase the efficacy of certain antibiotics such as ceftazidime and levofloxacin.

A study was done in 2011 by [23], and found that heartwood of *Acacia catechu* has anti-fungal activity. The obtained results demonstrated that the higher concentration of the extract showed good antifungal activity against all the fungal strains tested, and the lower concentration showed very weak or no activities.

Acacia catechu thus have various medical property [24]. Studies were done in 2010 to assess the antimicrobial activity of the *Acacia catechu* against organisms causing dental caries. Based on the results, *Acacia catechu* willd has been proven to be a potent antimicrobial agent against dental infections like dental caries being caused primarily by *Streptococcus mutans* which is similar to the current result.

Another study was done by [25] to evaluate the antibacterial activity of heartwood extract of *Acacia catechu* willd. The results demonstrated that the antibacterial activity of the extracts (Ethanolic and Aqueous) at different concentrations was determined by measuring the zone of inhibition. The ethanolic extract was more effective against *Staphylococcus aureus* with a zone of inhibition of 24 mm diameter (at conc200 g.) and was least effective against *Pseudomonas aeruginosa* and *Bacillus subtilis* with a zone of inhibition of 10mm (at conc. 200 g.) and 11mm (at conc. 200 g.) respectively. *E. coli* showed a zone of inhibition of 19mm diameter (at conc. 200 g.) and *Klebsiella pneumoniae* showed inhibition zone of 16mm diameter (at conc. 200 g.) among the other studied bacterial species. It was concluded that the *Acacia catechu* willd heart wood extracts contain antibacterial activity

[26, 27] performed a study on the Antimicrobial property of *Acacia* species. The results concluded that *A. nilotica* (pods) and *A. catechu* (bark) were reported to be the most active against different bacterial and fungal strains. The mathanolic extract of *A. nilotica* (pods) showed the highest activity against *E. coli*, *S. aureus* and *A. niger*, whereas *A. catechu* exhibited its prominent activity against *S. aureus* and *C. albicans*. Although, the hexane extract of *A. nilotica* was also found the most active against *S. typhi*. In whole antimicrobial experiment, *Acacia Jacquemonti* was found to have the weakest or no activity.

The increasing evidence has supported that the plants of genus *Acacia* contain high amount of bioactive secondary compound, and therefore, are likely to be promising in drug discovery. The importance of secondary compounds in *Acacia* lie in a variety of their functions, that the most important among them include their being anticancer (triterpenoid and saponins), diuretic (glucosides), natriuretic (glucosides), important nutraceutical (polysaccharide and gum) anti-digestive disorder (saponins, tannins and flavanoids), anti-oxidant (polyphenols), antiplasmodial (treptamine, tannins, organic acids and saponins) [25, 28].

As the above discussion has elaborated, *Acacia catechu* is of a high medical value. One such property is anti-microbial activity. Various studies reported the anti-bacterial activity of the plant. They showed a great response against *streptococcus* mutants and *lactobacillus acidophilus*. These two organisms play a major role in the dental caries. The anti-bacterial activity of *Acacia catechu* is positivity higher against *S. mutans* than the later *L. acidophilus*. *Acacia catechu* showed less activity on these organisms when compared to chlorhexidine, but the former is a herbal drug with no side effects.

CONCLUSION

Taxifolin isolated from *Acacia catechu* leaf was found to be effective as antibacterial against different bacterial pathogens, providing the scientific basis for its traditional application in Indian folk medicine against many Oro dental infections. Further studies should be done to find out the active compound responsible for antibacterial effects, and other necessary pharmacological studies are needed to be conducted to use it in modern drugs' developments.

REFERENCES

1. Karouche Saida, Khaldi Sofiane, Benbott Amel. (2018). Phytochemical, Free Radical Scavenging and Antimicrobial Activities of the Maize Stigmas, Collected of Ain Mlila (East Algeria). *World Journal of Environmental Biosciences*, 7 (4): 35-40.
2. Ali Al Ghasham, Mohammed Al Muzaini, Kamal Ahmad Qureshi, Gamal Osman Elhassan, Riyaz Ahmed Khan, Syeda Ayesha Farhana. (2017). Phytochemical Screening, Antioxidant and Antimicrobial Activities of Methanolic Extract of *Ziziphus mauritiana* Lam. Leaves Collected from Unaizah, Saudi Arabia. *International Journal of Pharmaceutical Research & Allied Sciences*, 6(3):33-46
3. Palombo, E.A. Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases.2008. *EvidBased Compl. Alt.*
4. Rishton, G.M. Natural products as a robust source of new drugs and drug leads: past successes and present day issues. *Am. J. Cardiol.*,2008, 101, 43D49D.
5. Sivak Elena, Bugaev Sergey, Sokolov Mikhail, Glinushkin Alexey. (2018). Antimicrobial bio-components form red algae species: a review of application and health benefits. *Entomology and Applied Science Letters*, 5 (3): 85-90.
6. Chemistry of Chinese yew, *Taxus chinensis* var. *mairei*. Cunfang Li, Changhong Huo, Manli Zhang, Qingwen Shi, *Biochemical Systematics and Ecology*, Volume 36, Issue 4, April 2008, Pages 266–282.
7. Rao PR, Seshadri TR, L-Epi-catechin from *Acacia catechu*, *Journal Scientist Indian Research*, 7B, 1948, 59.
8. Bibhabasu Hazra, (Bose Institute), Kolkata, Rhitajit Sarkar, Santanu Biswas, Nripendranath Mandal, The Antioxidant, Iron Chelating and DNA Protective Properties of 70% Methanolic Extract of 'Katha' (Heartwood extract of *Acacia catechu*) *journal of complementary & integrative medicine* vol.7 / 2010 /issue 1.
9. Rao PR, Seshadri TR, LEpicatechin from *Acacia catechu*, *Journal Scientist Indian Research*, 7B, 1948, 59.
10. Qadry JS, Shah's and Qadry's Pharmacognosy, 12th edition, B.S Shah Prakashan, Ahmedabad, 2008, 302303.
11. Ray D, Sharatchandra KH, Thokchom IS, Antipyretic, antidiarrhoeal, hypoglycaemic and Hepatoprotective activities of ethyl acetate extract of *Acacia catechu* Willd. In albino rats, *Indian Journal of Pharmacology*, 38(6), 2006, 408413.
12. Makena PS, Pierce SC, Chung KT, Sinclair SE. Comparative mutagenic effects of structurally similar flavonoids quercetin and taxifolin on tester strains *Salmonella typhimurium* TA102 and *Escherichia coli* WP-2uvrA'. *Environ Mol Mutagenesis* 2009; 50:451-9.
13. Whiley RA and Beighton D, "Streptococcus and oral streptococci" bite-sized tutorial. July 2013.
14. Ahrne, S, Nobaek, S, Jeppsson, B, Adlerberth, I, Wold, AE, Molin, G. 1998. The normal *Lactobacillus* flora of healthy human rectal and oral mucosa. *J Appl Microbiol.* 85(1):88–94.
15. Bunting RW, Crowley M, Hard DG, Keller M (1929). The prevention of dental caries through the limitation of growth of *Bacillus acidophilus* in the mouth. *J Am Dent Assoc*16:224-230.
16. Collins, CH and Lyne, P.M 1976. *Microbiological methods*, London, Butterworths and co.288p.
17. Betty A., Forbes. Daniel F. Sahm., Alice S. Weissfeld. Bailey & Scott's *Diagnostic Microbiology* 11th edition Mosby page 229 – 257.
18. Lawrence R. Antimicrobial activity of *Acacia catechu* bark extracts against selected pathogenic bacteria. *Int J Curr Microbiol Appl Sci* 2015; 1:213-22.

19. Kuspradini H, Mitsunaga T, Ohashi H. Antimicrobial activity against *Streptococcus sobrinus* and glucosyltransferase inhibitory activity of taxifolin and some flavanone rhamnosides from kempas (*Koompassia malaccensis*) extracts. *J Wood Sci* 2009; 55:308-13.
20. An SM, Kim HJ, Kim JE, Boo YC. Flavonoids, taxifolin and luteolin attenuate cellular melanogenesis despite increasing tyrosinase protein levels. *Phytother Res* 2008; 22:1200-7.
21. McDonald, M.; Dougall, A.; Holt, D.; Huygens, F.; Oppedisano, F.; Giffard, P.M.; Inman-
22. Bamber, J.; Stephens, A.J.; Towers, R.; Carapetis, J.R.; Currie, B.J. Use of a single nucleotide polymorphism genotyping system to demonstrate the unique epidemiology of methicillin-resistant *Staphylococcus aureus* in remote aboriginal communities. *J. Clin. Microbiol.* 2006, 44, 3720-3727.
23. In vitro evaluation of anti mycotic activity of heartwood extract of *Acacia catechu* WilldA Roy, RV Geetha, T Lakshmi - *Journal of pharmacy research*, 2011.
24. V. Gayathri Devi, Anitha John, R. Sreekala Devi, V. A. Prabhakaran. Pharmacognostical studies on *Acacia catechu* willd and identification of antioxidant principles. *International journal of pharmacy and pharmaceutical sciences*. Vol 3, suppl 2, 2011.
25. Role of botanicals as antimicrobial agents in management of dental infections—a review S Dhinahar T Lakshmi *Int. J. Pharm. Bio Sci*, 2011.
26. Patel, Jayshree¹; Kumar, Vipin¹; Bhatt, Shreyas² Antimicrobial screening and phytochemical analysis of the resin part of *Acacia catechu* *pharmaceutical biology journal* January 2009, Vol. 47, No. 1, Pages 3437.
27. Steiner M, Lin RS (June 1998). "Changes in platelet function and susceptibility of lipoproteins to oxidation associated with administration of aged garlic extract". *J Cardiovasc Pharmacol* 31 (6): 904–8.
28. Jayasekhar P, Mohanan PV, Hepatoprotective activity of ethyl acetate extract of *Acacia catechu*, *Indian Journal of Pharmacology*, 29(6), 1997, 426-428.