



Research Article

ISSN : 2277-3657
CODEN(USA) : IJPRPM

In Silico Prediction of SSRs and Functional Annotation of ESTs from Catharanthus Roseus

Pravej Alam*, Thamer Albalwai

Department of Biology, College of Science and Humanities, Prince Sattam bin Abdulaziz University, Al-Kharj 11942, Saudi Arabia

*Email: [alamprez @ gmail.com](mailto:alamprez@gmail.com)

ABSTRACT

Catharanthus roseus (periwinkles) belongs to the Apocynaceae family with great anti-cancer, anti-diabetic, and hepatoprotective values. Due to the large number of active molecules accumulated in these plants, they are of particular concern, especially in the pharmaceutical sector. The availability of ESTs gave the genetic algorithm of the plant to differentiate between the species accessions at the genetic level. The high-throughput method used for mining and detection of microsatellites (SSRs) embedded in ESTs gave a new insight for molecular markers' development. 19899 ESTs were retrieved, examined by NCBI EST dB and assembled 2692 to get full-length contigs sequences. 338 microsatellites (SSR) loci were predicted with an average of SSR per 9.33 kb of EST though MISA-web tools out of 2692 contigs. Furthermore, trinucleotide, a well-known SSR was examined and found to be the most favorable repeats' type (26.62%) followed by dinucleotide (24.22), mononucleotide (48.22%), and hexanucleotide (0.3%) types. The highest frequency of (A/T)*n* was reported in this finding followed by (AAG)*n*. The simple sequence repeats (SSR) extracted from *C. roseus* EST's data were used as molecular tools for genetic characterization in the present study. These predicted SSRs can be significantly used for constructing the genetic maps and also for differentiating the accession between the species.

Key words: *C. roseus*, ESTs, Contigs, SSR.

INTRODUCTION

Madagascar periwinkle (*Catharanthus roseus* L.) is a dicotyledonous and miracle medicinal plant with antitumor bioactive compounds [1-3]. The plant is domesticated and cultivated worldwide for ornamental and medicinal use [4, 5]. *C. roseus* is also one of the best sources for terpenoid indole alkaloids synthesized with a wide range of plant metabolites [6-8]. It has antidiabetic properties when the alcoholic extract of the plant is given to the streptozotocin-induced diabetic rats and shows a remarkable effect in lowering of glycemia in both diabetic and normal rats [9, 10]. Since many species have been evolved worldwide, it is difficult to segregate the plant at the species level.

Molecular markers (RAPD, AFLP, and SSR) are extensively used for analyses of genetic diversity, mapping of loci or gene, and marker-assisted selection in breeding technology [11-13]. These microsatellites or SSRs are distributed in plant genomes showing co-dominance and reproducibility. It has been frequently used for genetic scrutiny in breeding technology and it is recognized as a consistent and crucial method in plant genetics [14]. The advantages of SSRs development gave efficient hypervariability, codominant, and widespread genomic to differentiate the plant species at an inclusive spectrum [15].

METHODOLOGY

19899 *C. roseus* express sequence tags (ESTs) were retrieved from dB EST of NCBI for the analyses of SSRs. The EST sequences were passed through sequence cleaning and masking, through high throughput web application EGassembler [16] contrary to the NCBI *Vec* screen (<https://www.ncbi.nlm.nih.gov/tools/vecscreen/>) to remove the vector contaminant, poly A/T, and short sequences (adapter). The following parameters' minimum match (>10) and minimum score (20) with no stretch of (A/T)_n the existing step-wise process were applied in EGassembler.

Finally, in a non-redundant dataset of 2692 contigs, CAP3 program (EGassembler) was used to assemble the contigs for further analyses. MISA web (<https://webblast.ipk-gatersleben.de>), a web SSR finder developed by Beier et al. [17] was used to predict the EST-SSR loci.

Functional Annotations

The functional annotations with references to the biological process of assembled EST-contigs (2692) were predicted by the BLAST2go program obtained from Omics data. Based on NCBI BLAST and gene ontology, the biological function was predicted.

RESULTS AND DISCUSSION

A total of 19899 redundant ESTs' sequences were carried out for SSR analyses, retrieved from NCBI EST dB having 10196333 bp in *C. roseus* genome. During pre-processing, the sequence clean, masking of vector contaminant, low complexity sequences, and Poly A/T tails were examined and assembled effectively from 100087bp *C. roseus* ESTs. After mining of 19899 ESTs (5963 trimmed), 2692 contigs were finally generated to obtain the hypervariable class, I microsatellites i.e SSRs. The MISA web application (<https://webblast.ipk-gatersleben.de>) was used to screen the SSR through the mining program to search the 1-6 nucleotide repeat motifs. In this program, it was observed that only 338 hypervariable SSR loci were developed from 2692 contigs (Table 1). The SSR loci frequency was 9.33 kb per ESTs analyzed in *C. roseus*.

Table 1: The results of microsatellite search of *Catharanthus roseus* ESTs

Parameters	Values
Total number of ESTs	19899
Total sequence analyzed in bp	10196333
Total masking in bp	100087
EST after vector and Poly A/T removal	19868
Total number of singletons	5646
Total number of sequences examined:	2692
Total size of examined sequences (bp):	1911399
Total number of SSR loci located	338
Frequency of SSR loci in <i>Catharanthus roseus</i> EST	1 per 9.66 kb

The obtained frequency per SSR loci from *C. roseus* contigs was found satisfactory and in agreement with other findings in many plants (Figure 1). It is also noted that this finding is also in agreement with previous studies reported in rice (3.4), soybean (7.4), and maize (8.1) [18-20].

Repeats	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	Total
A	-	-	-	-	-	30	14	10	9	2	3	3	1	1		73
C	-	-	-	-	-		1			3						4
G	-	-	-	-	-	2	1	1				1				5
T	-	-	-	-	-	30	20	9	5	4	5	3	1	1	3	81
AG	-		3	3	2											8
AT	-	5	1	3												9
CT	-	8	3	4	4											19
GA	-	3		3												6
GT	-	1														1
TA	-	2	7	5	1											15
TC	-	7	8	7	3											25
TG	-	1														1
AAG	4	1														5
ACT						1										1
AGA	3															3
AGC		2														2
ATC	1	1														2
ATG	2															2
ATT	1															1
CAA	3	1														4
CAC	1															1
CAG	1	1														2
CAT	1	1														2
CCA		1														1
CCG		1														1
CCT		1														1
CGA		1														1
CGC	1															1
CGG	2															2
CTC	2															2
CTT	7															7
GAA	5	2														7
GAC	1															1
GAT	3	2														5
GCA	6	1														7
GCC	1	1														2
GCG		1														1
GCT		1														1
GGA		2														2
GGC		1														1
GTC	1	1														2
GTG	1															1
GTT	1															1
TAA		1														1
TAT	1															1
TCA	1	1														2
TCC	1															1
TCT	1															1
TGA	2															2
TGC	2	1														3
TTA	1															1
TTC	5	1														6
CAGCCA	1															1

Figure 1 : Frequency of identified SSR motifs from assembled ESTs of *C. roseus*

Repeats	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	Total
A/T	-	-	-	-	-	60	34	19	14	6	8	6	2	2	3	154
C/G	-	-	-	-	-	2	2	1		3		1				9
AC/GT	-	2														2
AG/CT	-	18	14	17	9											58
AT/AT	-	7	8	8	1											24
AAC/GTT	4	1														5
AAG/CTT	25	4														29
AAT/ATT	3	1														4
ACC/GGT	2	1														3
ACG/CGT	2	2														4
ACT/AGT						1										1
AGC/CTG	9	6														15
AGG/CCT	3	3														6
ATC/ATG	10	5														15
CCG/CGG	4	4														8
ACAGCC/CTGTGG	1															1

Figure 2 : Frequency of classified repeat types (considering sequence complementary) from assembled ESTs of *C. roseus*

The developed microsatellites or SSRs were characterized as the signature sequences in terms of simple motif type or compound motif or both. The obtained 338 SSRs from *C. roseus* are simple repeat motifs consisted of mononucleotide to hexanucleotide. The maximum frequency of motif was recorded as mononucleotide (48.22%) trailed by the trinucleotide (26.62%), dinucleotide (24.22), and hexanucleotide types (0.29%). The finding of trinucleotide repeat as the main single sequence repeats ; played an important role in previously reported plants (Figure 2) [21, 22]. The trinucleotide was earlier reported in wheat (32%), and sorghum (49%) with the signature sequence (CCG)_n. similarly, the (AAG)_n motif repeats have shown the maximum or rich motif in the trinucleotide repeat of *Gossypium barbadense* and *Curcuma longa*, which is in agreement with our findings (AAG)_n as shown in Table. 4 [23, 24]. The obtained dinucleotide (non-trimeric repeats) rather than trinucleotide SSRs may not be qualified as the best SSRs nomenclature due to some mutations [20]. The resulted SSR repeat motifs obtained from the ESTs-contigs were shown in Figure 3 as A/T (45.56%), AG/CT (17.16%), AAG/CTT (8.58%), AGC/CTG (4.44%), ATC/ATG (4.44%), CCG/CGG (2.37), and AGG/CTT (1.78%) rich frequency respectively. The furthest common repeat motifs were predicted as A/T, AG/CT, AAG/CTT, ACAGCC/CTGTGG with the frequency of 45.56%, 17.16%, 8.58%, 31.42%, 15.68% and 8.98% and 0.3%, respectively (Table 2). The biological function was also predicted by using the method of the Blast2Go program on the Omics box and they were recorded as GO:0008150 (biological process), GO:0050896 response to stimulus and GO:0051716 (cellular process) followed by others [25, 26] (Figure 4; Table 3).

Table 2: Maximum frequency of SSR repeats based on nucleotide repeats from assembled ESTs of *C. roseus*

Repeats based on nucleotide	SSRs repeats	% Frequency	Profuse motif	% Frequency
Mononucleotide	163	48.22	A/T	45.56
dinucleotide	84	24.22	AG/CT	17.16
Trinucleotide	90	26.62	AT/AT	7.10
Hexanucleotide	1	0.29	AAG/CTT	8.58
Total	338	-	AGC/CTG	4.44
-	-	-	ATC/ATG	4.44




Figure 3: Graphical representation of SSR-motif with their distribution frequency analyzed from assembled EST of *C. roseus*




Figure 4: Gene ontology-based functional annotation of EST-contigs obtained from *C. roseus* showing the biological functions

Table 3: Functional annotation of ESTs-contigs for biological process

Level	GO ID	GO Name	GO Type	Parents (ACC)	Parents (Name)
1	GO:0008150	Biological process	Biological Process	-	-
2	GO:0050896	Response to stimulus	Biological Process	GO:0008150	Biological process
2	GO:0009987	Cellular process	Biological Process	GO:0008150	Biological process
3	GO:0051716	Cellular response to stimulus	Biological Process	GO:0009987, GO:0050896	Cellular process, response to stimulus
3	GO:0042221	Response to chemical	Biological Process	GO:0050896	Response to stimulus
4	GO:0010035	Response to inorganic substance	Biological Process	GO:0042221	Response to chemical
4	GO:0001101	Response to acid chemical	Biological Process	GO:0042221	Response to chemical
4	GO:0070887	Cellular response to chemical stimulus	Biological Process	GO:0051716, GO:0042221	Cellular response to stimulus, response to chemical

5	GO:0071229	Cellular response to acid chemical	Biological Process	GO:0001101, GO:0070887	Response to acid chemical, cellular response to chemical stimulus
5	GO:1902617	Response to fluoride	Biological Process	GO:0010035, GO:0001101	Response to inorganic substance, response to acid chemical
6	GO:1902618	Cellular response to fluoride	Biological Process	GO:0071229, GO:1902617	Cellular response to acid chemical, response to fluoride

CONCLUSION

338 non-redundant hypervariable SSR type I obtained from EST of *C. roseus* using SSR predicted from MISA web tool is found to be reproducible, cost-effective, and time-saving. These 338 non-redundant SSR information may give a better platform to understand the genetic variation and help in the genome mapping of *C. roseus*. The functional annotation also provided information about ESTs involved in various processes. It may be used to provide the deep functional gene network information that is involved in particular metabolite synthesis with their deep functional annotations in *C. roseus* genome.

ACKNOWLEDGMENT

This Publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University.

REFERENCES

1. Alam P., Khan Z.A., Abdin, M.Z., Khan, J.A., Ahmad, P., Elkholy, S.F., Sharaf-Eldin, M.A., Efficient regeneration and improved sonication-assisted Agrobacterium transformation (SAAT) method for *Catharanthus roseus*. 3 Biotech. 2017, 7(1):26. doi: 10.1007/s13205-016-0593-5.
2. Pucot J R, Manting M M E, Demayo C G. Ethnobotanical Plants used by Selected Indigenous Peoples of Mindanao, the Philippines as Cancer Therapeutics. Pharmacophores. 2019; 10(3): 61-69.
3. Baranitharan M, Tamizhazhagan V, Koven-dan K. Medicinal Plants as Potent Power for Malaria Control: Review. Entomol. Appl. Sci. Lett. 2019; 6(1): 28-44.
4. Mujib, A., Ali, M., Isah, T., Dipti, Somatic embryo mediated mass production of *Catharanthus roseus* in culture vessel (bioreactor)—a comparative study. Saudi J Biol Sci. 2014, 21:442–449.
5. Medhini N, Divakara Y G, Prabha D, Manjulakumari D. Bioefficacy of *Calendula officinalis* Linn. (Asteraceae) extracts in the control of *Spodoptera litura* Fabricus (Noctuidae: Lepidoptera) under laboratory conditions. J. Biochem. Tech. 2012; 3(5): S167-S169.
6. Valdiani, A., Kadir, M. A., Tan, S. G. Talei, D., Abdullah, M. P., Nikzad, S., Nain-e havandi *Andrographis paniculata* present yesterday, absent today: a plenary review on underutilized herb of Iran's pharmaceutical plants, Mol. Biol. Rep. 2012, 39:5, 5409–5424
7. Facchini, P. J., Alkaloid biosynthesis in plants: biochemistry, cell biology, molecular regulation, and metabolic engineering applications, Annu. Rev. of Plant Biol, 2001, 52: 29–66
8. Roepke, J., Salim, V., Wu, M., Thamm, A.M., Murata, J., Ploss, K., Boland, W. and De Luca, V., Vinca drug components accumulate exclusively in leaf exudates of Madagascar periwinkle. PNAS, 2010, 107, 34, 15287-92.
9. Kanjekar A P. On Anti-Diabetic Potential of Phyto-nanoparticles Comparison with Hormonal Therapy and Medicinal Plants. Int. J. Pharm. Phytopharm. Res. 2019; 9(1): 103-111.
10. Wodu C O, Iwuji S C, Adienbo O M. Antihyperglycaemic activity of piper guineense in diabetic female albino wistar rats. Int. J. Pharm. Phytopharm. Res. 2017; 7(2): 1-4.
11. Park, Y.H., West, M.A.L., Clair, D.A.S., Evaluation of AFLPs for germplasm fingerprinting and assessment of genetic diversity in cultivars of tomato (*Lycopersicon esculentum* L.). Genome. 2004; 47: 510–518
12. Hend, B.T., Ghada, B., Sana, B.M., Mohamed, M., Mokhtar, T., Amel, S.H., Genetic relatedness among Tunisian plum cultivars by random amplified polymorphic DNA analysis and evaluation of phenotypic characters. Sci Hortic. 2009; 121: 440–446

13. Pirseyedi, S.M., Valizadehghan, S., Mardi, M., Ghaffari, M.R., Mahmoodi, P., Zahravi, M., Zeinalabedini, M. and Nekoui, S.M.K. Isolation and characterization of novel microsatellite markers in pomegranate (*Punica granatum L.*). *Int J Mol Sci.* 2010; 11: 2010–2016
14. Goldstein, D. B., and Schlötterer, C., Microsatellites: evolution and applications. *Q. Rev. Biol.* 1999. 83: 633–634
15. Agarwal, M., Shrivastava, N., Padh, H., Advances in molecular marker techniques and their applications in plant sciences. *Plant Cell Rep.* 2008; 27:617–631
16. Masoudi-Nejad, A., Tonomura, K., Kawashima, S., Moriya, Y., Suzuki, M., Itoh, M., Kanehisa, M., Endo, T., Goto, S., EGassembler: online bioinformatics service for large-scale processing, clustering and assembling ESTs and genomic DNA fragments. *Nucleic Acids Res.* 2006, 34:459-462.
17. Beier, S., Thiel, T., Münch, T., Scholz, U., Mascher, M., MISA-web: a web server for microsatellite prediction, *Bioinformatics*, 2017, 33, (16): 2583–2585
18. Cardle, L., Ramsay, L., Milbourne, D., Macaulay, M., Marshall, D., Waugh, R., Computational and experimental characterization of physically clustered simple sequence repeats in plants. *Genetics.* 2000, 156(2):847-54.
19. Gao, L., Tang, J., Li, H. and Jia, J. Analysis of microsatellites in major crops assessed by computational and experimental approaches. *Molecular Breeding* 12, 2003, 245–261
20. Joshi, R.K., Kuanar, A., Mohanty, S., Subudhi, E., Nayak, S., Mining and characterization of EST derived microsatellites in *Curcuma longa L.* *Bioinformation.* 2010; 20;5(3):128-31
21. Ramsay, L., Macaulay, M., degli Ivanissevich, S., MacLean, K., Cardle, L., Fuller, J., Edwards, K.J., Tuveesson, S., Morgante, M., Massari, A., Maestri, E., Marmioli, N., Sjakste, T., Ganal, M., Powell, W., Waugh, R., A simple sequence repeat-based linkage map of barley. *Genetics.* 2000 156(4):1997-2005
22. Edenilson, R., Adriane Nunes de, S., Saito, Daniel, Tsai, Mui S., In silico characterization of microsatellites in *Eucalyptus spp.*: abundance, length variation and transposon associations. *Genetics and Molecular Biology*, 2005, 28: 582-588
23. Kantety, R.V., La Rota, M., Matthews, D.E., Sorrells, M.E., Data mining for simple sequence repeats in expressed sequence tags from barley, maize, rice, sorghum and wheat. *Plant Mol Biol.* 2002: 48(5-6):501-10
24. Gupta, P.K., Balyan, H.S., Sharma, P.C., Ramesh, B., Microsatellites in plants : a new class of molecular markers *Current Science*, 1996, 70, 45-54
25. Cai, K., Zhu, L., Zhang, K., Li, L., Zhao, Z., Zeng, W., Lin, X., Development and Characterization of EST-SSR Markers From RNA-Seq Data in *Phyllostachys violascens*. *Front. Plant Sci.* 2019, 10:50. doi: 10.3389/fpls.2019.00050
26. Singh, S., Gupta, S., Mani, A., Chaturvedi, A., Mining and gene ontology based annotation of SSR markers from expressed sequence tags of *Humulus lupulus*. *Bioinformation.* 2012; 8(3):114-22.